

SEP



Enero 2022

Núm. 53

POLIBOTÁNICA



POLIBOTÁNICA

ISSN 1405-2768



Núm. 53

 CONACYT
Consejo Nacional de Ciencia y Tecnología

Enero 2022



PÁG.

CONTENIDO

- 1 Fabáceas del Área de Protección de Flora y Fauna Médanos de Samalayuca, Chihuahua, México
Fabaceas of the Flora and Fauna Protection Area Médanos de Samalayuca, Chihuahua, Mexico
J.R. Rueda-Torres | L. De León Pesqueira | A.B. Gatica Colima
- 13 Estudio de la flora presente en apíarios de tres municipios en el estado de Yucatán, México
Study of flora present in apiaries of three municipalities in the state of Yucatan, Mexico
C.I. Briceño Santiago | J. Cano Sosa | A.L. Ramos Díaz | R. Noriega Trejo | D.I. Couoh May
- 35 Relaciones filogenéticas de especies de *Phaseolus* de México con base en marcadores de ADN cloroplástico
Phylogenetic relationships of Phaseolus species from México based on chloroplastic DNA markers
V.H. Villarreal Villagrán | J. S. Muruaga Martínez | M.L.P. Vargas Vázquez | N. Mayek Pérez | S. Hernández Delgado
- 53 Las ingestiones e islas de los bosques espinosos del Caldenal dentro de los pastizales Austral Pampeanos
The Caldenal thorny forests ingestions and islands inside the Austral Pampean grasslands
E.L. Guerrero
- 69 Diversidad de especies de plantas arvenses en tres monocultivos del Bajío, México
Diversity of weed species in three monocultures from Bajío, Mexico
R. Guzmán Mendoza | V. Hernández Hernández | M.D. Salas Araiza | H.G. Núñez Palenius
- 87 Genetic diversity and genetic structure of *Capsicum annuum* L., from wild, homegarden and cultivated populations in a heterogeneous environment in Oaxaca, Mexico
Diversidad genética y estructura genética de Capsicum annuum L., de poblaciones silvestres, de traspatio y cultivadas en un ambiente heterogéneo en Oaxaca, México
R.T. Tapiaez | J.M. Peñaloza Ramírez | A.P. Olvera | A.L. Albaran Lara | K. Oyama
- 105 Morfología polínica de *Neomillspaughia* y *Podopterus* (Polygonaceae: Eriogonoideae: Coccolobeae)
Pollen morphology of Neomillspaughia and Podopterus (Polygonaceae: Eriogonoideae: Coccolobeae)
K.C. Durán Escalante | J.J. Ortiz Díaz | M. M. Ferrer | J. Tun Garrido
- 119 Morfoanatomía, histoquímica y germinación de las semillas de *Mammillaria parkinsonii* Ehrenb. (Cactaceae)
Morphoanatomy, histochemistry and germination of the seeds of Mammillaria parkinsonii Ehrenb. (Cactaceae)
Y. Uribe Salazar | A. Quintanar Isaias | C. Barbosa Martínez | J. Flores | C.L. Jiménez Sierra
- 135 Asymbiotic germination, *ex situ* conservation and *in vitro* plant regeneration of *Catasetum integerrimum* Hook
Germinación asimbiótica, conservación ex situ e in vitro regeneración de plantas de Catasetum integerrimum Hook
G. López Puc | G.J. Herrera Cool
- 151 Fitoquímicos y propiedades nutraceuticas de durazno (*Prunus persica* L.) cultivado en Zacatecas
*Phytochemicals and nutraceutical properties of peach (*Prunus persica* L.) harvested in Zacatecas*
J. Aguayo Rojas | S. Mora Rochín | X. Tovar Jiménez | J.J. Rochín Medina | R.O. Navarro Cortez
- 167 Evaluation of extracts of endemic trees (*Magnolia* spp.) in Mexico against the fruit fly pest and preliminary phytochemical study
*Evaluación de extractos de árboles endémicos (*Magnolia* spp.) de México contra la plaga de la mosca de la fruta y estudio fitoquímico preliminar*
S.G. Vásquez Morales | E.A. Alvarez Vega | D.A. Infante Rodríguez | J.P. Huchin Mian | M. Pedraza Reyes
- 183 Características fenotípicas, nutricionales y nutraceuticas de frutos de chile x'catik, dulce y su híbrido f1 (*Capsicum annuum* L.)
*Phenotypic, nutritional and nutraceutical traits of x'catik chili fruits, sweet and its f1 hybrid (*Capsicum annuum* L.)*
Y.A. Mis Valdez | M.J. Hernández Pinto | R. Garruña | K.B. Medina Dzul | R.H. Andueza Noh
- 197 Mecanismos de infección endógena en frutos de cacao con *Moniliophthora roreri*
Mechanisms of endogenous infection in cocoa fruits with Moniliophthora roreri
V. Flores | L. Gómez Rodríguez | J.A. López García | J. Grajales Conesa
- 211 Efectos de *Bacillus subtilis* cepas GBO3 y IN937b en el crecimiento de maíz (*Zea mays* L.)
*Effects of *Bacillus subtilis* strains GBO3 and IN937b on the growth of corn (*Zea mays* L.)*
A.E. Gutiérrez Calvo | A. Gutiérrez Estrada | C.L. Miceli Méndez | M.A. López Miceli
- 219 Conhecimentos etnobotânicos de mateiros em comunidades rurais da região da Serra das Almas, Paraná - Brasil
Conocimiento etnobotánico de los silvicultores en comunidades rurales de la región Serra das Almas, Paraná - Brasil
Ethnobotanical knowledge of foresters in rural communities in the Serra das Almas region, Paraná - Brazil
M. Ferreira Clarindo | A. Staniski | J. Strachulski
- 239 Valor cultural de los recursos forestales no maderables en comunidades zapotecas de la Sierra Juárez de Oaxaca
Cultural value of non-timber forest resources in Zapotec communities of the Sierra Juarez de Oaxaca
J. Martínez López | N.G. Molina Luna | S. Rangel Landa | C. Aquino Vázquez | A. Acosta Ramos



Portada

Podopterus mexicanus Humb. & Bonpl. Polygonaceae. “Rompe capa”. Árboles o arbustos de 1.5-6.0 m de altura, ramas con espina terminal, braquiblastos, hojas fasciculadas, flores blancas a verdosas en fascículos, y frutos cubiertos por el perianto externo que forma alas delgadas y largamente decurrentes hacia el pedicelo. Crece en bosques tropicales caducifolios y bosques espinosos, sobre suelo rocoso negro derivado de rocas ígneas. En elevaciones de 550-760 m. Florece de abril a mayo y fructifica de junio a septiembre. Se distribuye desde México hasta Centroamérica. En México se encuentra en los estados de Colima, Guerrero, Oaxaca, Puebla, Tamaulipas, Veracruz y Yucatán. Se utiliza como planta melífera y para leña.

Podopterus mexicanus Humb. & Bonpl. Polygonaceae. "Rompe capa". Trees or shrubs 1.5-6.0 m tall, branches with a terminal spine, brachyblasts, fasciculate leaves, white to greenish flowers in fascicles, and fruit covered by the external perianth that forms thin wings and longly decurrent towards the pedicel. It grows in tropical deciduous forests and thorny forests, on black, rocky soil derived from igneous rocks. At elevations of 550-760 m. Blossoms from April to May and bears fruit from June to September. It is distributed from Mexico to Central America. In Mexico it is found in the states of Colima, Guerrero, Oaxaca, Puebla, Tamaulipas, Veracruz and Yucatán. It is used as a melliferous plant and for firewood.

por/by Rafael Fernández Nava



INSTITUTO POLITÉCNICO NACIONAL

Director General: *Dr. Arturo Reyes Sandoval*

Secretario General: *Mtro. Juan Manuel Cantú Vázquez*

Secretario Académico: *Dr. David Jaramillo Vigueras*

Secretario de Extensión e Integración Social: *Dr. Luis Alfonso Villa Vargas*

Secretario de Investigación y Posgrado: *Dra. Laura Arreola Mendoza*

Secretario de Servicios Educativos: *Dra. Ana Lilia Coria Páez*

Secretario de Administración: *M. en C. Javier Tapia Santoyo*

Director de Educación Superior: *Mtro. Mauricio Igor Jasso Zaranda*

ESCUELA NACIONAL DE CIENCIAS BIOLÓGICAS

Directora:

Dra. Yadira Rivera Espinoza

Subdirectora Académica:

M. en C. Martha Patricia Cervantes Cervantes

Subdirector Administrativo:

Ing. Raúl Chávez Alvircio

Jefe de la Sección de Estudios de Posgrado e Investigación:

Dr. Gerardo Aparicio Ozores

Subdirector de Servicios Educativos e Integración Social:

Dr. Felipe Neri Rodríguez Casasola

POLIBOTÁNICA, Año 27, No. 53, enero-junio 2022, es una publicación semestral editada por el Instituto Politécnico Nacional, a través de la Escuela Nacional de Ciencias Biológicas. Unidad Profesional Lázaro Cárdenas, Prolongación de Carpio y Plan de Ayala s/n, Col. Santo Tomás C.P. 11340 Delegación Miguel Hidalgo México, D.F. Teléfono 57296000 ext. 62331.

<http://www.herbario.encb.ipn.mx/>, Editor responsable: Rafael Fernández Nava. Reserva de Derechos al Uso Exclusivo del Título No. 04-2015-011309001300-203. ISSN impreso: 1405-2768, ISSN digital: 2395-9525, ambos otorgados por el Instituto Nacional del Derecho de Autor. Responsable de la última actualización de este número, Unidad de informática de la ENCB del IPN, Rafael Fernández Nava, Unidad Profesional Lázaro Cárdenas, Prolongación de Carpio y Plan de Ayala s/n, Col. Santo Tomás C.P. 11340 Delegación Miguel Hidalgo México, D.F.

Las opiniones expresadas por los autores no necesariamente reflejan la postura del editor de la publicación.

Queda estrictamente prohibida la reproducción total o parcial de los contenidos e imágenes de la publicación sin previa autorización del Instituto Politécnico Nacional.

REVISTA BOTÁNICA INTERNACIONAL DEL INSTITUTO POLITÉCNICO NACIONAL

EDITOR EN JEFE
Rafael Fernández Nava

EDITORAS ASOCIADAS
Maria de la Luz Arreguín Sánchez

COMITÉ EDITORIAL INTERNACIONAL

Christiane Anderson
University of Michigan
Ann Arbor. Michigan, US

Edith V. Gómez Sosa
Instituto de Botánica Darwinion
Buenos Aires, Argentina

Heike Vibrans
Colegio de Postgraduados
Estado de México, México

Jorge Llorente Bousquets
Universidad Nacional Autónoma de México
Ciudad de México, México

Graciela Calderón de Rzedowski
Instituto de Ecología del Bajío
Pátzcuaro, Mich., México

Delia Fernández González
Universidad de León
León, España

Theodore S. Cochran
University of Wisconsin
Madison, Wisconsin, US

Jerzy Rzedowski Rotter
Instituto de Ecología del Bajío
Pátzcuaro, Mich., México

Hugo Cota Sánchez
University of Saskatchewan
Saskatoon, Saskatchewan, Canada

Luis Gerardo Zepeda Vallejo
Instituto Politécnico Nacional
Ciudad de México, México

Fernando Chiang Cabrera
Universidad Nacional Autónoma de México
Ciudad de México, México

Claude Sastre
Muséum National d'Histoire Naturelle
París, Francia

Thomas F. Daniel
California Academy of Sciences
San Francisco, California, US

Mauricio Velayos Rodríguez
Real Jardín Botánico
Madrid, España

Francisco de Asís Dos Santos
Universidad Estadual de Feira de Santana
Feira de Santana, Brasil

Noemí Waksman de Torres
Universidad Autónoma de Nuevo León
Monterrey, NL, México

Carlos Fabián Vargas Mendoza
Instituto Politécnico Nacional
Ciudad de México, México

Julieta Carranza Velázquez
Universidad de Costa Rica
San Pedro, Costa Rica

José Luis Godínez Ortega
Universidad Nacional Autónoma de México
Ciudad de México, México

Tom Wendt
University of Texas
Austin, Texas, US

José Manuel Rico Ordaz
Universidad de Oviedo
Oviedo, España

DISEÑO Y FORMACIÓN ELECTRÓNICA
Luz Elena Tejeda Hernández

OPEN JOURNAL SYSTEM Y TECNOLOGÍAS DE LA INFORMACIÓN
Pedro Aráoz Palomino

Toda correspondencia relacionada con la revista deberá ser dirigida a:

Dr. Rafael Fernández Nava

Editor en Jefe de

POLIBOTÁNICA

Departamento de Botánica

Escuela Nacional de Ciencias Biológicas, Instituto Politécnico Nacional
Apdo. Postal 17-564, CP 11410, Ciudad de México

Correo electrónico:

polibotanica@gmail.com

rfernán@ipn.mx

Dirección Web

http://www.polibotanica.mx

POLIBOTÁNICA es una revista indexada en:

CONACYT, índice de Revistas Mexicanas de Investigación Científica y Tecnológica del Consejo Nacional de Ciencia y Tecnología.

SciELO - Scientific Electronic Library Online.

Google Académico - Google Scholar.

DOAJ, Directorio de Revistas de Acceso Público.

Dialnet portal de difusión de la producción científica hispana.

REDIB Red Iberoamericana de Innovación y Conocimiento Científico.

LATINDEX, Sistema regional de información en línea para revistas científicas de América Latina, el Caribe, España y Portugal.

PERIODICA, Indice de Revistas Latinoamericanas en Ciencias.



Polibotánica

ISSN electrónico: 2395-9525

polibotanica@gmail.com

Instituto Politécnico Nacional

México

<http://www.polibotanica.mx>

**GENETIC DIVERSITY AND GENETIC
STRUCTURE OF *Capsicum annuum* L.,
FROM WILD, BACKYARD AND
CULTIVATED POPULATIONS IN A
HETEROGENEOUS ENVIRONMENT IN
OAXACA, MEXICO**

**DIVERSIDAD GENÉTICA Y ESTRUCTURA
GENÉTICA DE *Capsicum annuum* L., DE
POBLACIONES SILVESTRES, DE
HUERTOS FAMILIARES Y
CULTIVADAS EN UN AMBIENTE
HETEROGÉNEO EN OAXACA, MÉXICO**

Toxqui-Tapia, R.; J.M. Peñaloza-Ramírez; A. Pacheco-Olvera; A.L. Albarran-Lara and K. Oyama.
GENETIC DIVERSITY AND GENETIC STRUCTURE OF *Capsicum annuum* L., FROM
WILD, BACKYARD AND CULTIVATED POPULATIONS IN A HETEROGENEOUS
ENVIRONMENT IN OAXACA, MEXICO.

DIVERSIDAD GENÉTICA Y ESTRUCTURA GENÉTICA DE *Capsicum annuum* L., DE
POBLACIONES SILVESTRES, DE HUERTOS FAMILIARES Y CULTIVADAS EN UN
AMBIENTE HETEROGÉNEO EN OAXACA, MÉXICO.



GENETIC DIVERSITY AND GENETIC STRUCTURE OF *Capsicum annuum* L., FROM WILD, BACKYARD AND CULTIVATED POPULATIONS IN A HETEROGENEOUS ENVIRONMENT IN OAXACA, MEXICO.

DIVERSIDAD GENÉTICA Y ESTRUCTURA GENÉTICA DE *Capsicum annuum* L., DE POBLACIONES SILVESTRES, DE HUERTOS FAMILIARES Y CULTIVADAS EN UN AMBIENTE HETEROGÉNEO EN OAXACA, MÉXICO.

Toxqui-Tapia, R.;
J.M. Peñaloza-Ramírez;
A. Pacheco-Olvera;
A.L. Albaran-Lara
& K. Oyama.

GENETIC DIVERSITY AND GENETIC STRUCTURE OF *Capsicum annuum* L., FROM WILD, BACKYARD AND CULTIVATED POPULATIONS IN A HETEROGENEOUS ENVIRONMENT IN OAXACA, MEXICO.

DIVERSIDAD GENÉTICA Y ESTRUCTURA GENÉTICA DE *Capsicum annuum* L., DE POBLACIONES SILVESTRES, DE HUERTOS FAMILIARES Y CULTIVADAS EN UN AMBIENTE HETEROGÉNEO EN OAXACA, MÉXICO.

R. Toxqui-Tapia

*Escuela Nacional de Estudios Superiores Unidad Morelia,
Universidad Nacional Autónoma de México (UNAM).
Antigua Carretera a Pátzcuaro No. 8701, Col. Ex-Hacienda de San José de la
Huerta, 58190, Morelia, México (Phone: +52 (443) 322-27-04).*

J. M. Peñaloza-Ramírez / jmpenalo@uacam.mx

*Centro de Estudios de Desarrolló Sustentable y Aprovechamiento de la vida
Silvestre, CEDESU, Universidad Autónoma de Campeche, UAC. Campus 6, Avenida
Héroe de Nacozari #480, San Francisco de Campeche, México*

A. Pacheco-Olvera

*Facultad de Agronomía, Universidad Autónoma de Sinaloa (UAS),
Carr. Culiacán-El Dorado Km 17.5. Tel. y Fax 01 (667) 846-1084.
Apdo. Postal 726. Culiacán Sinaloa, México.*

L. Albaran-Lara

K. Oyama

*Escuela Nacional de Estudios Superiores Unidad Morelia,
Universidad Nacional Autónoma de México (UNAM).
Antigua Carretera a Pátzcuaro No. 8701, Col. Ex-Hacienda de San José de la
Huerta, 58190, Morelia, México (Phone: +52 (443) 322-27-04).*

RESUMEN: Las especies domesticadas han sido expuestas a presiones selectivas que regulan la estructura genética y divergencia entre poblaciones. Estimamos la diversidad genética, la estructura, la endogamia, el tamaño efectivo de la población, los cuellos de botella poblacionales y la divergencia del nicho ecológico, de poblaciones silvestres, de huertos familiares y cultivadas de *C. annuum*. Analizamos, 13 poblaciones en Oaxaca, México y usando siete marcadores de microsatélites nucleares. Las poblaciones de *C. annuum*, mostraron niveles altos de diversidad genética en poblaciones silvestres, de huertos familiares y cultivadas. Los chiles domesticados, en hábitats agrícolas tradicionales, contienen variedades genéticas únicas y por lo tanto, actúan como reservorios de variación genética y conservan la biodiversidad. Que indica, una fuerte selección, que ocurre en diferentes fenotipos lo que podría generar variedades genéticamente diversos, lo que ha favorecido la adaptación de las poblaciones a su entorno y ambientes. Los valores de la endogamia y el tamaño efectivo de la población son suficientes para permitir el mantenimiento de las poblaciones. Es necesario iniciar programas de selección de variedades de chile, para rescatar algunas características del fruto. La diferenciación genética, es moderada entre poblaciones, encontramos cuatro regiones geográficas que son muy divergentes entre ellas. Sugerimos, que en las poblaciones de chile, una fuerte selección artificial podría generar variedades genéticamente divergentes. Postulamos que la complejidad climática y los gradientes altitudinales en el estado de Oaxaca podrían promover la biodiversidad, la domesticación y la diversificación. Los chiles

Oaxaca son un ejemplo de diversificación de las razas, que incluyen chiles como, 'Tusta', 'Tabaquero', 'Solterito', 'Piquín', 'Nanchita', 'Costeño' and 'Chile de Agua'.

Palabras clave: *C. annuum*, proceso de domesticación, diversidad genética, cuellos de botella poblacionales, tamaño efectivo de la población.

ABSTRACT: Domesticated species have been exposed to artificial pressures that regulate the genetic structure and divergence between populations. We estimated the genetic diversity, structure, inbreeding, effective population size, population bottlenecks and the ecological niche divergence, of wild, homegarden and cultivated populations of *C. annuum*. We analyzed, 13 populations across Oaxaca, Mexico and using seven nuclear microsatellite markers. *C. annuum*, shown high levels of genetic diversity in wild, homegarden and cultivated populations. Domesticated chiles in traditional agricultural habitats harbor unique gene pools and thus act as genetic variation reservoirs and to preserve its biodiversity. That indicates, a strong selection occurred at different phenotypes, could generating genetically diverse materials that have favored population adaptation to locations and environments. The values of inbreeding and the effective population size were sufficient to allow the maintenance of the populations. It is necessary to initiate programs to select chile varieties, to rescue some traits in the fruit. Genetic differentiation, was moderated among populations, we detected four geographic regions that are highly divergent between them. We suggest that in chile populations, a strong artificial selection could generate genetically divergent materials. We postulate, that climatic complexity and altitudinal gradients in Oaxaca, could promoting biodiversity, domestication and diversification. The Chiles from Oaxaca, are an example to the diversification of landraces that include chiles such as, 'Tusta', 'Tabaquero', 'Solterito', 'Piquín', 'Nanchita', 'Costeño' and 'Chile de Agua'.

Key words: *C. annuum*, domestication process, genetic diversity, population bottlenecks, effective population size.

INTRODUCTION

Domesticated species have been exposed during their evolution to artificial selection that determine the genetic structure, phenotypic patterns and divergence among cultivated and wild populations (Charlesworth *et al.*, 1997; Pickersgill, 2007). Also, artificial selection operates in the domestication processes and has encouraged morphological, physiological, and genetic modifications that could alter mating systems and genetic structure in domesticated organisms (Renaut & Rieseberg, 2015). Some of the consequences of domestication, include loss of genetic diversity, introduction of these genes associated with domestication process, an accumulation of deleterious variants, phenotype convergence and local adaptation to plantations and could represent a profound impact on genomes (Meyer & Purugganan, 2013; Larson *et al.*, 2014).

Particularly, an important crop species in Mexico, the species *C. annuum* L. (Solanaceae) is known as chile and represented one of the most important crop, with a great economic significance (Pickersgill, 2007). The domestication process of *C. annuum* was conducted in Mexico, due to the morphological variation of varieties (López-Espinosa *et al.*, 2018). Chile represents, one of the ancestral crops of Mexico, where domestication process began between 5000-7000-BC, and occurred at the Tehuacán Valley (Meyer & Purugganan, 2013). Most of local varieties, have been cultivated for long periods with an intermediate cultivation and an agricultural management system (González *et al.*, 2011; Guzman *et al.*, 2019). Regarding their economic importance, several studies evaluated the importance of local varieties of chile using morphological, cytogenetic and genetic markers (Contreras-Toledo *et al.*, 2011). These studies shown, that chile is one of the primary crop in America and Mexico and that different species were domesticated in

several provinces independently (Ibiza *et al.*, 2012). Nevertheless, further studies are needed to estimate heterozygosity levels to implement conservation programs for the rational use of the remaining diversity of wild populations (Rodríguez *et al.*, 2007; Guzman *et al.*, 2019).

Oaxaca state, is considered as a region with a high diversity of species (Ortiz-Pérez *et al.*, 2004), as well as many local domesticated species (Contreras-Toledo *et al.*, 2011). Oaxaca possess a high biological and cultural diversity, that coupled with a complex environmental heterogeneity, results in their high biodiversity, represented by more than 12 500 species of flora and fauna; many of which have been utilized by local populations (Contreras-Toledo *et al.*, 2011). An example, of native chilies domesticated at the Oaxaca state, 'Tusta', 'Tabaquero', 'Solterito', 'Piquín', 'Nanchita', 'Costeño' and 'Chile de Agua' (Alonso *et al.*, 2008; Vera-Guzmán *et al.*, 2011). That represented a different degree of domestication, artificial selection and genetic differentiation.

Taking into account the economic importance of this chile resource, is necessary to evaluate their genetic structure to generate procedures for management and conservation. The main objectives of this study are to examine genetic diversity levels of populations of *C. annuum* with different levels of human management located in the valleys and mountains in Oaxaca, Mexico. We addressed the following questions: 1) We characterized, the population genetic diversity, differentiation and genetic inbreeding levels, to test if genetic diversity parameters could be affected by the level of disturbance. (2) We evaluated, the possible occurrence of recent bottlenecks events, estimated the effective population size and the niche similarity and divergence. To understand whether populations undergone recent bottleneck processes and to recognize if populations located in different valleys and mountains significantly differentiated from each other.

MATERIALS AND METHODS

Study system

C. annuum plants are recognized for their bright coloured, fleshy, podlike fruit. Plants are bushy and low growing 30-38cm tall. The stems are to some extent woody with thin, green leaves 4-10cm long and 1-4cm wide (González *et al.*, 2011). White flowers are produced from leaf axils in summer, but are inconspicuous. The fruit usually remains ornamental for 8 to 12 weeks. This species has been organized into five botanical groups of which only three cherry, cone and cluster peppers are familiar potted plants (Pickersgill, 2007). Botanical classification of *Capsicum* has been difficult, due to the high number of varieties, the deficiency of well-defined traits and the presence of hybridization identified of some species (Ibiza *et al.*, 2012). Chile represented, one of the ancestral and utilized crops of Mexico and domestication occurred at the Tehuacán Valley. Most of the local crops, have been cultivated and consumed for long time with a stable cultivated under an agricultural management system that constitute an important genetic resource (Guzman *et al.*, 2019).

Collecting sites

The study area was designated among two biogeographic regions at the Oaxaca State in Mexico. Young leaves were collected, five from wild populations, five from homegarden and three from domesticated populations at the Oaxaca State (Table 1, Fig. 2). We collected 13 populations distributed in Oaxaca and sampled between 12 - 20 individuals. We sampled from trees separated with at least 1000 m. For sampling, we choose the criteria of geographic location. Genomic DNA was extracted from 100 mg of fresh leaf material using the protocol designed by Lefort & Douglas, (1999). Seven nuclear microsatellite loci, were amplified using the loci previously reported for *C.*

annuum (Minamiyama *et al.*, 2006) and utilizing multiplex polymerase chain reaction approach. PCR techniques were arranged into three different groups. The first group include the primer pairs Agi021, Agi069, Agi098 and Agi111. The second group comprise the primers CAMS163, CAMS460 and CAMS679 (Minamiyama *et al.*, 2006). PCR techniques were performed using the QIAGEN Multiplex PCR kit (QIAGEN) in a 5 μ l volume containing 1X Multiplex PCR Master Mix, 2 μ M of primer, dH₂O, and 20 ng of template DNA. The thermal cycling conditions consisted of 35 cycles of 94°C for 1 min, annealing 51°C for 1 min, extension at 72°C for 2 min and final extension at 72°C for 10 min. Multiplex PCR products were combined with a GeneScan-500 LIZ size standard and performed on an ABI-PRISM 3100 Avant sequencer (Applied Biosystems). Fragments were analyzed and recorded using the Peak Scanner program 1.0 (Applied Biosystems).

Table 1. Locality name, sample size, geographical coordinates, mean number of alleles (A), mean observed heterozygosity (H_O), mean expected heterozygosity (H_E), for 13 populations of *C. annuum* in the Oaxaca, Mexico, separated by wild, homegarden and cultivated populations. Standard errors are included in parenthesis.

Locality	Sample size	Genetic diversity			
		Coordinates	A	H_O	H_E
Wild populations					
1. El Coyul1	11	15.413/ -97.420	3.71 (2.07)	0.700 (0.10)	0.556 (0.08)
2. Zaachila1	7	18.619/ -97.548	5.00 (2.78)	0.707 (0.24)	0.671 (0.17)
3. Proterill	10	18.403/ -97.548	6.14 (2.53)	0.728 (0.11)	0.716 (0.14)
4. Guleagui	13	18.703/ -97.603	4.28 (3.03)	0.698 (0.12)	0.589 (0.10)
5. MorroM2	11	18.555/ -97.635	4.14 (2.08)	0.589 (0.08)	0.568 (0.07)
Total			4.654	0.684	0.620
Homegarden					
6. Agua de Soll	11	18.865/ -97.702	4.71 (2.31)	0.771 (0.16)	0.576 (0.10)
7. Agua de Sol2	11	18.464/ -97.568	4.42 (2.66)	0.696 (0.08)	0.560 (0.08)
8. Zaachila2	7	18.409/ -97.429	4.28 (0.98)	0.833 (0.16)	0.658 (0.10)
9. MorroM3	10	18.358/ -97.480	3.57 (2.07)	0.757 (0.10)	0.532 (0.07)
10. El Coyul2	11	18.476/ -97.572	4.57 (1.21)	0.650 (0.10)	0.536 (0.03)
Total			4.31	0.741	0.572
Cultivated					
11. Taviche1	10	18.619/ -97.477	4.71 (2.40)	0.620 (0.12)	0.665 (0.07)
12. Taviche2	11	18.605/ -97.542	5.14 (1.94)	0.767 (0.21)	0.698 (0.10)
13. Taviche3	11	18.636/ -97.422	5.57 (1.37)	0.712 (0.14)	0.639 (0.03)
Total			5.14	0.699	0.667

Genetic diversity analysis

For each *C. annuum* populations, the following genetic diversity parameters: number of alleles per locus (A), observed heterozygosity (H_O), expected heterozygosity (H_E) and inbreeding coefficient (F_{IS}), using GENETIX 4.02 software (Belkhir *et al.*, 1996).

Genetic structure and Bayesian admixture analysis

To test genetic structure, we performed a hierarchical test of population structure (AMOVA), employing the stepwise mutation model (SMM) using ARLEQUIN 3.5. program (Excoffier, & Lischer, 2010). We grouped, *C. annuum* populations taking into account the domestication degree: wild, homegarden and cultivated populations that represented a different extent of domestication, artificial selection and genetic differentiation. Statistical significance was tested using 10^4 permutations using ARLEQUIN program. We estimated the gene flow between *C. annuum* populations was

considered by estimating the Bayesian-scaled long-term effective population size (N_e) and migration rate (m) employing the MIGRATE 3.5.1 software (Beerli & Felsenstein, 2001). For the analyses, the starting chain value was set to 20^6 visited and 16 recorded genealogies, following a burn-in period of 50^3 iterations.

The genetic ancestry of each individual plant of *C. annum* was analyzed with the STRUCTURE 2.3.4 software (Pritchard *et al.*, 2000; Falush *et al.*, 2003; Hubisz *et al.*, 2009). STRUCTURE uses a Bayesian clustering model to determine the proportion of each individual's ancestry originating from different populations (Evanno *et al.*, 2005). The optimal number (K) of groups was determined by varying K from 1 to 10 (e.g. to achieve the effective number of groups) and running the analysis ten times for each K value to find the maximum posterior likelihood [$\text{LnP}(\text{D})$]. Each run was performed using 10^6 Markov chain Monte Carlo (MCMC) repetitions following a burn-in period of 50^4 iterations. We used an admixture model that allows the correlation of allele frequencies without any a priori information. Following the procedure of Evanno *et al.* (2005), we determined the most likely value of K based on the maximum value of ΔK .

We utilized, two different approaches to achieve genetic clustering, with the programs STRUCTURE and ADEGENET, to contrast the differences among genetic groups. We utilized the Discriminant Analysis of Principal Components (DAPC) using the adegenet 2.0 (Jombart *et al.*, 2010) package in R Version 3.5.1. We implemented the 'find.clusters' function to achieve the discriminant analysis of principal component procedure (Jombart *et al.*, 2010). To avoid overfitting, 1000 principal components (e.g. to accomplish the correct number of clusters) were retained, accounting for ~70% of the total variance. This analysis runs successive K-means clustering and the optimal number of K was selected based on the lowest associated Bayesian Information Criteria (BIC) value after examining the rate of decrease in BIC. We tested values of genetic groups $K = 1 - 20$ (e.g. to achieve correctly the number of genetic groups) with 10 replicates of each K . The DAPC function was executed using this clustering, retaining 2 axes of principal components analysis (PCA) sufficient to explain 80% of the total variance of the data (Jombart *et al.*, 2010).

To identify geographic barriers and genetic discontinuity among *C. annum* populations, we used the Monmonier's maximum difference algorithm with BARRIER v. 2.2 software (Manni *et al.*, 2004). BARRIER creates a map of the sampling localities from their geographical coordinates. Barriers are then represented on the map by identifying the maximum values within the population-pairwise genetic distance matrix. We used a pairwise matrix of average square distances (ASD) estimated for *C. annum* populations. Resampling random subsets of individuals within populations estimated with the MSA program with 100 bootstrap replicate distances that were used to achieve statistical significance for the predicted barriers.

Testing for niche divergence and conservatism

To determine the ecological niche differentiation between genetic groups *C. annum* and identify the suitable habitat between genetic groups, we performed niche divergence test, niche overlap, range overlap and niche similarity test using 11 climate variables and using Schoener's D and Hellinger's I niche similarity metrics, implemented in ENMtools 1.4.4 software (Warren *et al.*, 2008) and the package ecospat (Broennimann *et al.*, 2012). These indexes quantify the niche similarity, which ranges from 0 (no overlap in habitat suitability) to 1 (identical niche models in habitat suitability). We applied a threshold in which habitat is considered suitable, using the average of between genetic groups "Minimum training Presence logistic" threshold (=0.20). Niches were considered different if the observed value of niche overlap was less than the niche overlap value from 95 or all 100 of the niche overlap values. Niche breadth was

calculated as proportional similarity between the observed distribution of the environmental variables (Warren *et al.*, 2008).

To analyze and visualize the ENMs of *C. annum*, we quantified the niche overlapping, niche equivalence (e.g. assess whether the ENMs between genetic groups are different than expected even if they share the same underlying distribution) and niche similarity (e.g. inquires whether one species niche can predict the occurrence of the other), estimated between genetic groups. For each *C. annum* genetic groups a PCA-Environmental approach derived from multivariate theory implemented in the package ecospat (Broennimann *et al.*, 2012). The procedure involves a principal component analysis (PCA) to associate climatic values with species occurrence densities in the current period (Warren *et al.*, 2008). All analyses were conducted using R program V. 3.2.3 and ecospat package (Broennimann *et al.*, 2012).

Changes in population size

We used the BOTTLENECK 1.2 software (Piry *et al.*, 1999) to detect population bottlenecks, using the domestication degree: wild, homegarden and cultivated populations of *C. annum*. Recent population bottlenecks that could be defined as a population where the rare alleles are the first to be lost decreasing the mean number of alleles per locus. In contrast, heterozygosity is less affected, producing a transient heterozygosity excess relative to that expected based on the resulting number of alleles (Luikart & Cornuet, 1998). For this test, we used 90% stepwise and 10% multistep mutations and performed 10^4 iterations, employing the Wilcoxon's signed-rank test, the stepwise mutation (SMM), the infinite allele (IAM) and the two-phase mutation (TPM) models. To contrast, the differences among the different mutation models. We estimated the effective population size (N_e) of all locations, using the LDNe software (Waples & Do, 2008). This program implemented the bias-correction technique developed by Waples & Do, (2008) to obtain N_e from a sample of S individuals. We set $P_{crit} = 0.02$ (i.e., alleles with a frequency < 0.02 are excluded), which generally provides a good balance between accuracy and bias. Confidence intervals (CIs) for N_e were calculated with the chi-square approximation implemented in LDNe.

RESULTS

Genetic diversity analysis

Values of genetic diversity parameters for *C. annum* ranging from high for homegarden locations ($A = 4.28\text{-}4.71$, $H_O = 0.699\text{-}0.833$) followed by moderate-high for wild locations ($A = 3.71\text{-}6.14$, $H_O = 0.589\text{-}0.728$) and finally lower values for cultivated locations ($A = 4.71\text{-}5.57$, $H_O = 0.620\text{-}0.767$) see Table 1. Wright's inbreeding coefficient within populations (F_{IS}) revealed in general, positive values representing a heterozygote "reduction" (Table 1); F_{IS} values were low for wild populations ($F_{IS} = 0.014\text{-}0.244$), low in homegarden populations ($F_{IS} = 0.170\text{-}0.279$) and cultivated populations ($F_{IS} = 0.032\text{-}0.121$) all values are significant see Table 1.

Genetic structure and Bayesian admixture analysis

Results from hierarchical analysis of molecular variance (AMOVA) separated by domestication degree of *C. annum* (Table 2), Indicated that most of the genetic variation resided within populations (75.95%), while the differentiation among groups only accounted for the remaining variation (22.76%). Followed by variation among populations within groups (1.28%), all levels are significant (Table 2). Genetic exchange results detected with MIGRATE (Table 3, Fig. 2) outlined that in agreement with genetic structuring, we detected less exchange among populations across long distance such the Cluster 2 (green), (i.e. Structure Analysis) have low exchange with respect to all populations. The rest of the populations that belong to Cluster 1 (blue), 3 (yellow) and 4

(red) show clear indications of genetic exchange values ranging from low to moderated values of exchange among the population and despite belonging to wild, homegarden and homegarden populations (Table 3, Fig. 2).

Table 2. Analysis of molecular variance (AMOVA) performed on the nSSR data and using RST. We grouped, the populations taking into account the domestication degree: wild, homegarden and cultivated populations that represented a different extent of domestication and genetic differentiation. Asterisks indicate statistically significant values ($P < 0.01$). Tests were based on 10^4 random permutations.

Source of variation	SS	Variance components	Percentage of variation	Fixation index
R_{ST}				
Among groups	2	0.037	1.28	$\Phi_{CT} = 0.090^{***}$
Among populations within groups	10	0.664	22.76	$\Phi_{SC} = 0.010^{***}$
Within populations	223	2.218	75.95	$\Phi_{ST} = 0.080^{***}$
Total	235.5	2.92104		

Table 3. Levels of genetic exchange estimated with the program MIGRATE. Directional pairwise migration rates among eleven *C. annuum* populations from Oaxaca Mexico. Donating and receiving populations are below and above the diagonal, respectively. Migration rates are given as the value of effective number of migrants per generation (Nm). Bold number indicate statistically significant values ($P < 0.01$).

<i>Nm</i>	MM2S1	GuelS2	Coy1T3	Ag1T4	Tav3C5	Tav2C6	MM3T7	ZacS8	Coy2Si9	Tav1C10	PotrS11	Za2T12	Ag2T13
MM2S1	—	0.145	0.423	1.034	0.065	1.065	3.432	3.055	1.845	0.525	0.460	0.188	0.756
GuelS2	0.705	—	0.398	1.791	1.099	5.860	0.787	1.094	1.589	1.733	1.736	0.099	9.047
Coy1T3	0.225	0.246	—	1.970	0.062	0.185	0.345	0.308	0.303	0.123	1.233	1.585	0.789
Ag1T4	2.195	2.191	1.788	—	1.676	1.806	1.366	1.378	1.254	2.162	2.193	0.448	2.070
Tav3C5	1.519	1.826	1.456	1.819	—	2.952	1.833	1.190	1.846	2.271	2.575	1.592	2.138
Tav2C6	2.247	2.045	1.879	2.879	2.498	—	2.475	2.625	1.978	1.574	2.482	2.062	2.802
MM3T7	2.722	0.741	0.635	0.635	0.635	2.137	—	2.725	2.550	0.212	1.745	2.751	1.587
ZacS8	2.732	0.287	0.431	2.909	2.160	2.915	2.481	—	1.747	1.193	1.341	0.436	1.183
Coy2Si9	3.470	1.194	1.614	2.614	2.475	1.732	2.236	2.128	—	2.833	2.310	1.300	2.200
Tav1C10	2.641	2.931	2.300	1.343	2.950	2.003	2.146	2.952	2.470	—	2.565	2.647	2.891
PotrS11	2.273	2.100	2.591	2.591	2.436	1.903	1.631	2.451	2.578	2.756	—	1.478	2.478
Za2T12	1.777	2.887	2.990	0.888	2.443	2.762	3.546	2.330	3.542	2.665	1.665	—	2.994
Ag2T13	0.872	2.378	2.865	2.865	2.702	2.084	0.996	0.872	2.084	0.996	0.872	2.084	—

Results, from the Bayesian clustering analysis showed that the maximum posterior likelihood [$\text{LnP}(D)$] and the maximum ΔK value showed that $K=4$ is the effective number of genetic groups (Fig. 1, Fig. 2). Also, the DAPC analysis (Fig. 3) shown the same clustering and corroborates the grouping detected with STRUCTURE. Clusters detected by the DAPC in Fig. 2 and 3 show the distribution of ancestry proportions in each collection site. Cluster 1 (blue) is moderately widespread across the Oaxaca basin and includes the Guelaguchi (wild), Potrerillo (wild) and el Coyul (wild) populations. Cluster 2 (green) is mostly restricted to the Oaxaca mountains and includes populations of Tavicche2 (cultivated), Tavicche3 (cultivated), Tavicche1 (cultivated), Agua de Sol2 (homegarden) and Agua de Sol1 (homegarden) which are

consistently structured across the landscape. Cluster 3 (yellow) is restricted to the Oaxaca basin, and includes the populations of El Coyul (homegarden) and Zaachilac2 (homegarden). Cluster 4 (red) has a moderated geographic distribution through the Oaxaca basin and includes the populations of Zaachilac1 (wild), Moro Mazatlan3 (homegarden) and Morro Mazatlan2 (wild) 1. Accordingly with the great structure shown that the clusters are very different from each other, the presence of admixture in the populations was scarcely evident (Fig. 2).




Fig. 1. Values of ΔK plotted against K , the peak indicates the most probable number of genetic groups given the data using Structure Harvester (Earl, 2011).




Fig. 2 Each pie chart represents the proportions in each population of the four genetic groups as assigned by the program STRUCTURE. Green, red, yellow and blue Genotype are representing the genetic ancestry groups corresponding to *C. annuum* populations. Genetic discontinuities (bold lines B-1–2) obtained with Monmonier's maximum difference algorithm on genetic distances derived from microsatellite allele frequencies.

DPAPC1 Discriminant function, *C. annum*




Fig. 3. Discriminant Analysis of Principal Component, ordination based on 13 *C. annum* populations. Inference of population structure as identified through DAPC. The corresponding BIC plot showing the optimal number of clusters ($K = 4$) and the DA eigenvalues for the displayed scatterplot.

The results from genetic and geographic barriers analysis between 13 populations of *C. annum*, revealed two barriers (with over 50% bootstrap support) (Fig. 2). The most significant barrier, with 95% bootstrap support, sets apart the populations of Taviche2 (cultivated), Taciche3 (cultivated), Taviche1 (cultivated), Agua de Sol2 (homegarden) and Agua de Soll (homegarden), located in the northwest part of the basin from the rest of the *C. annum* populations. The second barrier, with 89% bootstrap support, is a complex break separating the southwestern part from the central part of the Oaxaca basin. This indicates that, in agreement with the high structure found, some populations have become isolated, such as the populations El Coyul (wild) from el Coyul (homegarden), Zaachila1 (homegarden) and Potrerillo (wild) (Fig. 2).

Testing for niche divergence and conservatism

C. annum genetic groups, had significantly non-equivalent ENMs (Fig. 4). The niche identity test indicated a high degree of climatic niche differences with values ranging from D statistic 0.774 to 0.837 genetic groups. The niche breadth measurements supported the hypothesis that broader niches exist for both *C. annum*: genetic groups 80%; 79%. These results indicate that the *C. annum*: genetic groups occupied different climate zones indicated by the niche between genetic groups (Fig. 4). The overlap index was a low: 0.225 between *C. annum*: genetic groups with low to moderate values of spatial range overlap ($D = 0.261$, $I = 0.544$) and indicated the amount of climate that share among the genetic groups in their distribution (Fig. 4).




Fig. 4. Niche overlap densities plots using the PCA-Env scripts ecospat (Broennimann *et al* 2012) between environmental ranges of *C. annum*, for the different genetic groups (i.e. STRUCTURE) darker shading indicates higher density of occurrences of the species; solid contour lines, 100% of available environment; dashed contour lines, 50% of most common background environment. The available environment in the study areas are defined and by green and red lines when comparing the two species at the same time from the sympatric with populations with the same background area. The correlation circle shows the loadings of individual environmental variables to the two PCA axes and the contribution of each variables to the construction of the PCA-Env. Sympatry.

Changes in population size

Results from the analyses to detect evidence of recent bottlenecks (excess heterozygosity) using the infinite allele (IAM), the two-phase (TPM) and the stepwise mutation (SMM) models see Table 4. No significant results were found with the IAM, TPM and SMM models as an indication of no bottlenecks in the wild, homegarden and domesticated populations of *C. annum*. Results from the test to estimate the effective population size (N_e) for *C. annum*

populations, grouped according to domestication degree, shown that cultivated populations ($Ne = 96.2$ individuals) had the highest value followed by the wild group populations ($Ne = 33.6$ individuals) and homegarden populations ($Ne = 26.3$ individuals). All Ne estimates had a high Jackknife support and a good confidence interval (CIs) (see Table 4).

Table 4. Bottleneck analysis for *C. annuum* populations in the Oaxaca, Mexico using Wilcoxon rank test under infinite allele, stepwise mutation and two phase model. Parameters for TPM: variance = 10%, proportion of SMM=90%, estimation based on 104 replications. P, probability. IAM, infinite allele model; TPM, two phase model; SMM, stepwise mutation model. ** Indicate significant deviation from equilibrium as value less than 0.05. The results obtained, for the estimation of the population effective size for the three genetic groups, values obtained with the program LDNe.

Models	Wild	Homegarden	Cultivated
IAM	0.01953	0.00781	0.71094
TPM	0.99219	0.40625	0.98828
SMM	1.00000	0.96094	0.99609
LDNe	33.6	26.3	96.2

DISCUSSION

Domesticated species, have been exposed during their evolution to hundreds of years of human selection in multiple environments and cultural contexts. This leads to phenotypic diversity of fruits and seeds (Charlesworth *et al.*, 1997). Native, homegarden and wild landraces could be affected by numerous factors such as, ecological characteristics, mountainous systems, anthropogenic disturbances and artificial selection (Larson *et al.*, 2014; Renaut & Rieseberg, 2015).

A high genetic variation observed, was observed for *C. annuum* at the homegarden ($H_O = 0.833$), whereas wild ($H_O = 0.728$) and cultivated ($H_O = 0.620$) showed lower values. Oaxaca state represents, a biodiversity hotspot in Mexico (Ferrusquía-Villafranca, 1993; García-Mendoza, 2004) and possess a great number of domesticated species and genetic diversity in regional chiles (Corona-Torres *et al.*, 2000; Latournerie *et al.*, 2001). A possible explanation of this diversity could be linked to the variability of soils, mountainous systems and the management of native and cultivated plants by farmers (Aguilar-Rincón *et al.*, 2010; Pacheco-Olvera *et al.*, 2012; López-Espinosa *et al.*, 2018). Evaluation of genetic variability among the *Capsicum* genotypes result essential for selection. Assessment of such specific alleles and genotypes should be important, with respect to adaptability of crops, with respect to climate change and biotic and abiotic stresses (Pickersgill, 2007; Zizumbo-Villarreal & Colunga-García, 2010).

Genetic diversity reduction, has been observed in crops and is due to population bottlenecks and artificial selection (Charlesworth *et al.*, 1997). In our *C. annuum* populations, we detected low inbreeding and population size for wild ($F_{IS} = 0.244$, $Ne = 33.6$), homegarden ($F_{IS} = 0.170$, $Ne = 26.3$) and cultivated ($F_{IS} = 0.121$, $Ne = 96.2$) populations. We observed, for *C. annuum* populations, a slightly reduction in inbreeding, effective population size and not significant signals of genetic bottlenecks (Larson *et al.*, 2014; Renaut & Rieseberg, 2015). Domestication syndrome, signified the morphological traits shared by domestic, but not by wild ancestors. That could modified, reproductive traits such as, altered timing of flowering and altered color compounds. This syndrome, could affecting in isolated populations with self-fertilization, where little or no pollen flow is present (Contreras-Toledo *et al.*, 2011).

We suggested for *C. annuum* the future tendency for landraces that will be exposed to the loss of diversity, affecting the ability to respond to climate changes (Meyer & Purugganan, 2013). Also, fire rates and habitat fragmentation has led chile plantations, wild, homegarden and cultivated to be isolated in patches (Ibiza *et al.*, 2012). Depletion of genetic diversity, has been reported to increase susceptibility to infections, caused by fungi, (e.g. *Phytophthora* and *Verticillium*) (González *et al.*, 2011). Crop wild relatives, contain genes to improve and resilience. *Capsicum* plants produced capsaicinoids that reduces the growth of fungal pathogens in wild chile, *Capsicum chacoense* (Rodríguez *et al.*, 2007). Genetically modified crops, are associated with various risks, such as, introduction of genes to susceptible crops, which leads to extinction of wild populations (Hanski & Mononen, 2011). Conservation programs, are necessary directed to select wild chile. Also gene flow, could contribute to rescuing genetic diversity of local varieties, that can counteract inbreeding and disease susceptibility (Corona-Torres *et al.*, 2000).

Genetic structure and Bayesian admixture analysis

Our study found that, genetic structuring of *C. annuum* among wild, homegarden and domesticated chiles, is associated with geographic features in the range of chile at the Oaxaca (Aguilar-Rincón *et al.*, 2010; Pacheco-Olvera *et al.*, 2012).

Oaxaca state represented, a biodiversity, cultural and ecosystems hotspot (Ferrusquía-Villafranca, 1993). Also harbored, a complex heterogeneity of soils, climatic that are delimited on a mountainous ranges coursing in a NNW-SSE direction and separated by valleys with tropical deciduous and xerophytic elements in the central portion (Ortiz-Pérez *et al.*, 2004). These intricated, patterns resulted in a geographical barrier, contributing to isolation and genetic differentiation (García-Mendoza, 2004). We observed a great genetic diversity and differentiation that indicated, that domestication had an effect on the distribution of variation among domesticated chiles. Native chiles from Oaxaca, are an example of diversification of landraces used by humans such as, 'Tusta', 'Tabaquero', 'Solterito', 'Piquín', 'Nanchita', 'Costeño' and 'Chile de Agua' (Vera-Guzmán *et al.*, 2011). Mountainous landscapes in Oaxaca, are constituted by heterogeneous topography and environmental ranges (Hanski & Mononen, 2011). Steep valleys and mountain demarcate the plant population habitats where gene flow is reduced thus, favoring differentiation (Latournerie *et al.*, 2001). Genetic differentiation, could be incremented by drift associated with reduced population size (Pacheco-Olvera *et al.*, 2012). Artificial selection, could be acting on plant populations determining morphological, physiological, reproductive, and genetic changes, leading to phenotypic and genotypic divergence between wild and managed populations (Kwon *et al.*, 2005; Casas *et al.*, 2007). As was observed, for local landraces of chile, which have been consumed the wild and domesticated chile across Oaxaca state (Zizumbo-Villarreal & Colunga-García, 2010).

The Cluster 1 (blue), Cluster 3 (yellow) and Cluster 4 (red) (e.g. Structure analysis) are located on the Pacific coast is associated with the Barrier 2 (Fig. 2). This clear structuring, have been revealed with populations with little gene flow. In the Sierra Madre del Sur, the valley ranges at elevations from 1000 m, but toward south elevations rise to 1800 m and host xerophytic vegetation and acts as a barrier to gene flow among tropical deciduous and temperate forest (Ortiz-Pérez *et al.*, 2004). We suggest, that higher variation in habitat and climatic conditions combined with high elevations and geographic distances could explain genetic divergence (Larson *et al.*, 2014). Populations in the valleys, are geographically isolated from highlands, thus gene flow is reduced and increasing differentiation. We indicated that in *C. annuum* populations, a strong artificial selection could generate genetically diverse materials that has favored population adaptation to locations and environments (Hanski & Mononen, 2011; López-Espinosa *et al.*, 2018). That could generate, diverse and divergent landraces adapted to different agroecological and cultural environments (Kwon *et al.*, 2005; Zizumbo-Villarreal & Colunga-García, 2010). All those phenotypes producing fruit with the most desirable attributes according to local people are cultivated and represented the highest level of artificial selection (Alonso *et al.*, 2008; Casas *et al.*, 2007; Aguilar-Rincón *et al.*, 2010).

CONCLUSIONS

High genetic diversity for *C. annuum* at the Oaxaca state, that denoted a region with a great number of domesticated and richness in Mexico that could encouraging the domestication syndrome. Native chiles from Oaxaca, are an example of diversification of landraces used by human such as, 'Tusta', 'Tabaquero', 'Solterito', 'Piquín', 'Nanchita', 'Costeño' and 'Chile de Agua'. We indicate, that the degree of heterozygosity, allelic richness and private alleles could be used for improving. Evaluation of such genotypes should be important, with respect to adaptability of crops, with respect to climate change.

Low genetic inbreeding, for wild and homegarden and low effective population size for wild and cultivated. It is necessary, programs to rescuing chile varieties. Through In-situ, preserving economically important plants. Through Ex-situ, using Botanical Gardens. Also, gene flow, between communities could contributed to rescuing genetic diversity which can counteract inbreeding effects and disease susceptibility.

Genetic differentiation was moderate, for wild, homegarden and cultivated populations that underlined a clear genetic structuring. We suggest that for *C. annuum* populations, a strong artificial selection, could generated genetically diverse materials that stimulated adaptation to locations and environments. Domestication or selection, had an effect on genetic and phenotypic variation. Oaxacan chile landraces, are in the stage of adaptation to different agroecological and cultural environments. As was observed, for local landraces of chile, which have been consumed, wild and domesticated at the Oaxaca state.

LITERATURE CITED

- Aguilar-Rincón, V., Corona-Torres, H., López-López, L., Latournerie-Moreno, M., Ramírez-Meraz, H., Villalón-Mendoza, M., & Aguilar-Castillo, J. (2010). *Los chiles de México y su Distribución*. INIFAP, IT-Conkal, UANL, UAN, Montecillo, Texcoco, Estado de México., Estado de México., México.: SINAREFI, Colegio de Postgraduados.
- Alonso, R., Moya, C., Cabrera, A., Pilar, P., Quiroga, R., Rosales, M., & Zuart, J. (2008). Evaluación *in situ* de la variabilidad genética de los chiles silvestres (*Capsicum* spp.) en la región frailesca del estado de Chiapas, México. *Cultivos Tropicales*, 49-55. <http://www.redalyc.org/articulo.oa?>
- Belkhir, K., Borsig, P., Chikhi, L., Raufaste, N., & Bonhomme, F. (1996). *GENETIX 4.05, logiciel sous Windows TM pour la génétique des populations*. Laboratoire Génome, Populations, Interactions, Université de Montpellier II: CNRS UMR 5171. <http://www.genetix.univ-montp2.fr/genetix/genetix>
- Beerli, P., & Felsenstein, J. (2001). Maximum likelihood estimation of a migration matrix and effective population sizes in n subpopulations by using a coalescent approach. *Proceedings of the National Academy of Sciences of the United States of America*, 4563–4568. doi:<https://doi.org/10.1073/pnas.081068098>
- Broennimann, O., Fitzpatrick, M., Pearman, P., Petitpierre, B., Pellissier, L., Yoccoz, N., Thuiller, W., Fortin, M. J., Randin, C., Zimmermann, N. E., Graham, C. H., & Guisan, A. (2012). Measuring ecological niche overlap from occurrence and spatial. *Global Ecology and Biogeography*, 481–497. doi:<https://doi.org/10.1111/j.1466-8238.2011.00698.x>
- Casas, A., Otero-Arnaiz, A., Pérez-Negrón, E., & Valiente-Banuet, A. (2007). *In situ Management and Domestication of Plants in Mesoamerica*. *Annals of Botany*, 1101–1115. doi:<https://doi.org/10.1093/aob/mcm126>
- Charlesworth, B., Nordborg, M., & Charlesworth, B. (1997). The effects of local selection, balanced polymorphism and background selection on equilibrium patterns of genetic diversity in subdivided populations. *Genetics Research*, 155-174. doi:<https://doi.org/10.1017/S0016672397002954>

- Contreras-Tolelo, A., López, S., Santacruz, V., Valadez, M., Aguilar, R., Corona, T., & López, P. (2011). Diversidad genética en México de variedades nativas de chile 'poblano' mediante microsatélites. *Revista Fitotecnia Mexicana*, 225 – 232. doi:http://www.scielo.org.mx/scielo.php?script=sci_arttext&pid=S0187-73802011000400003
- BIBLIOGRAPHY VI 2058**
- Corona-Torres, T., García-Velázquez, A., Castillo-González, F., Montero-Tavera, V., & Azpiroz-Rivero, H. (2000). Caracterización isoenzimática de la diversidad genética de colectas de chile (*Capsicum annuum* L. y *Capsicum chinense* Jacq.). *Revista Chapingo Serie Horticultura*, 5-17. doi:<https://doi.org/10.35196/rfm.2018.3.227-236>
- Evanno, G., Regnaut, S., & Goudet, S. (2005). Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Molecular Ecology*, 2611–2620. doi:<https://doi.org/10.1111/j.1365-294X.2005.02553.x>
- Excoffier, L., & Lischer, H. (2010). Arlequin suite ver 3.5: A new series of programs to perform population genetics analyses under Linux and Windows. *Molecular Ecology Resources*, 564-567. doi:<https://doi.org/10.1111/j.1755-0998.2010.02847.x>
- Falush, D., Stephens, M., & Pritchard, J. (2003). Inference of population structure using multilocus genotype data: linked loci and correlated allele frequencies. *Genetics*, 1567–1587. doi:<https://doi.org/10.1093/genetics/164.4.1567>
- Ferrusquía-Villafranca, L. (1993). *Biological diversity of Mexico*. Oxford UK: Oxford University Press. <https://ci.nii.ac.jp/naid/10027450251/>
- García-Mendoza, A. J. (2004). *Integración del conocimiento florístico del estado*. (García-Mendoza, A. Ordoñez, M., & Briones-Salas, M. Edits.) *Biodiversidad de Oaxaca* (pp. 305–325) México, D.F. Instituto de Biología, México: UNAM-Fondo Oaxaqueño para la Conservación de la Naturaleza, World Wildlife Fund
- BIBLIOGRAPHY VI 2058**
- González, I., Arias, Y., Quiñones, M., Miranda, I., Rodríguez, Y., & Peteira, B. (2011). Variabilidad molecular de genotipos de pimiento (*Capsicum annuum* L.) del programa de mejoramiento genético para la resistencia a pvy. *Protección Vegetal*, 69-72. http://scielo.sld.cu/scielo.php?script=sci_arttext&pid=S1010-27522011000200001&lng=pt&nrm=iso&tlang=es
- Guzman, F. A., Moore, S., de Vicente, M., & Jahn, M. (2019). Microsatellites to enhance characterization, conservation and breeding value of *Capsicum* germplasm. *Genet Resour Crop Evol*, 569-585. doi:<https://doi.org/10.1007/s10722-019-00801-w>
- Hanski, I., & Mononen, T. (2011). Eco-evolutionary dynamics of dispersal in spatially heterogeneous environments. *Ecology Letters*, 1025–1034. doi:<https://doi.org/10.1111/j.1461-0248.2011.01671.x>
- Hubisz, M., Falush, D., Stephens, M., & Pritchard, J. (2009). Inferring weak population structure with the assistance of sample group information. *Molecular Ecology Resources*, 1322–1332. doi:<https://doi.org/10.1111/j.1755-0998.2009.02591.x>
- Ibiza, V. P., Blanca, J., Cañizares, J., & Nuez, F. (2012). Taxonomy and genetic diversity of domesticated *Capsicum* species in the Andean region. *Genetic Resources Crops and Evolution*, 1077–1088. doi:<https://doi.org/10.1007/s10722-011-9744-z>
- Jombart, T., Devillard, S., & Balloux, X. (2010). Discriminant analysis of principal components: a new method for the analysis of genetically structured populations. *BMC Genomics*, 11-94. doi:<https://doi.org/10.1186/1471-2156-11-94>
- Kwon, Y. S., Lee, J., Yi, G., Yi, S., Kim, K., Soh, E., Bae, K. M., Park, E. K., Song, I. H., & Kim, B. (2005). Use of SSR markers to complement tests of distinctiveness, uniformity, and stability (DUS) of pepper (*Capsicum annuum* L.) varieties. *Molecular Cells*, 428–435. doi:<https://pubmed.ncbi.nlm.nih.gov/15995361/>
- Larson, G., Piperno, D., Allaby, R., Purugganan, M., Andersson, L., Arollo-Kalin, M., Barton, L., Vigueira, C. C., Denham, T., Dobney, K., Doust, A. N., Gepts, P., Gilbert, M. T., Gremillion, K. J., Lucas, L., Lukens, L., Marshall, F. B., Olsen, K. M., Pires, J. C., Richerson, P. J., Rubio de Casas, R., Sanjur, O. I., Thomas, M. G., & Fuller, M. (2014). Current perspectives and the future of domestication studies. *Proceedings of the National Academy of Sciences of the United States of America*, 6139-6146. doi:<https://doi.org/10.1086/285580>

- Latournerie-Moreno, L., López-Vázquez, J. S., Castañón-Nájera, G., Mijangos-Cortéz, O., Espadas-Villamil, G., Pérez-Gutiérrez, A., & Ruiz-Sánchez, E. (2015). Evaluación agronómica de chile habanero (*Capsicum chinense* Jacq.). *Agroproductividad*, 24-29. <http://cicy.repositorioinstitucional.mx/jspui/handle/1003/498>
- Lefort, F., & Douglas, G. (1999). An efficient micro-method of DNA isolation from mature leaves of four hardwood tree species *Acer*, *Fraxinus*, *Prunus* and *Quercus*. *Annals of Forest Science*, 259-263. doi:<http://dx.doi.org/10.1051/forest:19990308>
- López-Espinosa, S. T., Latournerie-Moreno, L., Castañón-Nájera, G., Ruiz-Sánchez, G., Gómez-Leyva, J., Andueza-Noh, H., & Mijangos-Cortés, J. (2018). Genetic Diversity of habanero pepper (*Capsicum chinense* Jacq.) using ISSR. *Rev. Fitotec. Mex*, 227 - 236. doi:<https://doi.org/10.35196/rfm.2018.3.227-236>
- Luikart, G., & Cornuet, J. (1998). Empirical evaluation of a test for identifying recently bottlenecked populations from allele frequency data. *Biological Conservation*, 228-237. doi:<https://doi.org/10.1111/j.1523-1739.1998.96388.x>
- Manni, F., Guerard, E., & Heyer, E. (2004). Geographic patterns of (genetic, morphologic, linguistic) variation: how barriers can be detected by “Monmonier’s algorithm”. *Human Biology*, 173-190. doi:<https://doi.org/10.1353/hub.2004.0034>
- Meyer, R. S., & Purugganan, M. (2013). Evolution of crop species: genetics of domestication and diversification. *Nature Reviews Genetics*, 840-852. doi:<https://doi.org/10.1038/nrg3605>
- Minamiyama, Y., Masato, T., & Masashi, H. (2006). An SSR-based linkage map of *Capsicum annuum*. *Molecular Breeding*, 157-169. doi:<https://doi.org/10.1007/s11032-006-9024-0>
- Ortiz-Pérez, M. A., Hernández-Santana, J., & Figueroa-Mah-Eng, J. (2004). Reconocimiento fisiográfico y geomorfológico. (García-Mendoza, A.J., Ordoñez, M. J., & Briones-Salas, M. Eds.) *Biodiversidad de Oaxaca*, Instituto de Biología, UNAM México: Fondo Oaxaqueño para la Conservación de la NaturalezaWorld Wildlife Fund, México.
- Pacheco-Olvera, A., Hernández, V., Rocha, R., González, R., & Oyama, K. (2012). Genetic Diversity and Structure of Pepper (*Capsicum annuum* L.) from Northwestern Mexico Analyzed by Microsatellite Markers. *Crop Science*, 231-241. doi:<https://doi.org/10.2135/cropsci2011.06.0319>
- Pickersgill, B. (2007). Domestication of Plants in the Americas: Insights from Mendelian and Molecular Genetics. *Annals of Botany*, 925-940. doi:<https://doi.org/10.1093/aob/mcm193>
- Piry, S., Luikart, G., & Cornuet, J. (1999). BOTTLENECK:a computer program for detecting recent reductions in the effective population size using allele frequency data. *J Hered*, 502-503. doi:<https://doi.org/10.1093/jhered/90.4.502>
- Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using multilocus genotype data: linked loci and correlated allele frequencies. *Genetics*, 1567-1587. doi:<https://doi.org/10.1093/genetics/164.4.1567>
- BIBLIOGRAPHY VI
2058**
- Renaut, S., & Rieseberg, L. (2015). The accumulation of deleterious mutation as a consequence of domestication and improvement in sunflowers and other *Compositae* crops. *Molecular Biology and Evolution*, 2273-2283. doi:<https://doi.org/10.1007/10.1093/molbev/msv106>
- Rodríguez, J., Peña, O., Gil, M., Martínez, C., Manzo, F., & Salazar, L. (2007). Rescate in situ del chile “poblano” en Puebla, México. *Rev. Fitotec. Mex*, 25-32. doi:www.scielo.org.mx/scielo.php?script=sci_nlinks&pid=S2007-0934201600050100500021&lng=en
- Vera-Guzmán, A. M., Chávez-Servia, J., Carrillo-Rodríguez, J., & López, M. (2011). Phytochemical evaluation of wild and cultivated pepper (*Capsicum annuum* L. and *C. pubescens* Ruiz & Pav.) from Oaxaca, Mexico. *Chilean J. Agricultural Research*. doi:<http://dx.doi.org/10.4067/S0718-58392011000400013>

Recibido:
31/marzo/2021

Aceptado:
10/diciembre/2021

- Waples, R. S., & Do, C. (2008). LdNe: a program for estimating effective population size from data on linkage disequilibrium. *Molecular Ecology Resources*, 753-756. doi:<https://doi.org/10.1111/j.1755-0998.2007.02061.x>
- Warren, D. L., Glor, R., & Turelli, M. (2008). Environmental niche equivalency versus conservatism: quantitative approaches to niche evolution. *Evolution*, 2868–2883. doi:<https://doi.org/10.1111/j.1558-5646.2008.00482.x>
- Zizumbo-Villarreal, D., & Colunga-GarcíaMarín, P. (2010). Origin of agriculture and plant domestication in West Mesoamerica. *Genetics Resources and Crops Evolution*, 813–825. doi:<https://doi.org/10.1007/s10722-009-9521-4>