

SEP



Enero 2022

Núm. 53

POLIBOTÁNICA



POLIBOTÁNICA

ISSN 1405-2768



Núm. 53

 CONACYT
Consejo Nacional de Ciencia y Tecnología

Enero 2022



PÁG.

CONTENIDO

- 1 Fabáceas del Área de Protección de Flora y Fauna Médanos de Samalayuca, Chihuahua, México
Fabaceas of the Flora and Fauna Protection Area Médanos de Samalayuca, Chihuahua, Mexico
J.R. Rueda-Torres | L. De León Pesqueira | A.B. Gatica Colima
- 13 Estudio de la flora presente en apíarios de tres municipios en el estado de Yucatán, México
Study of flora present in apiaries of three municipalities in the state of Yucatan, Mexico
C.I. Briceño Santiago | J. Cano Sosa | A.L. Ramos Díaz | R. Noriega Trejo | D.I. Couoh May
- 35 Relaciones filogenéticas de especies de *Phaseolus* de México con base en marcadores de ADN cloroplástico
Phylogenetic relationships of Phaseolus species from México based on chloroplastic DNA markers
V.H. Villarreal Villagrán | J. S. Muruaga Martínez | M.L.P. Vargas Vázquez | N. Mayek Pérez | S. Hernández Delgado
- 53 Las ingestiones e islas de los bosques espinosos del Caldenal dentro de los pastizales Austral Pampeanos
The Caldenal thorny forests ingestions and islands inside the Austral Pampean grasslands
E.L. Guerrero
- 69 Diversidad de especies de plantas arvenses en tres monocultivos del Bajío, México
Diversity of weed species in three monocultures from Bajío, Mexico
R. Guzmán Mendoza | V. Hernández Hernández | M.D. Salas Araiza | H.G. Núñez Palenius
- 87 Genetic diversity and genetic structure of *Capsicum annuum* L., from wild, homegarden and cultivated populations in a heterogeneous environment in Oaxaca, Mexico
Diversidad genética y estructura genética de Capsicum annuum L., de poblaciones silvestres, de traspatio y cultivadas en un ambiente heterogéneo en Oaxaca, México
R.T. Tapiaez | J.M. Peñaloza Ramírez | A.P. Olvera | A.L. Albaran Lara | K. Oyama
- 105 Morfología polínica de *Neomillspaughia* y *Podopterus* (Polygonaceae: Eriogonoideae: Coccolobeae)
Pollen morphology of Neomillspaughia and Podopterus (Polygonaceae: Eriogonoideae: Coccolobeae)
K.C. Durán Escalante | J.J. Ortiz Díaz | M. M. Ferrer | J. Tun Garrido
- 119 Morfoanatomía, histoquímica y germinación de las semillas de *Mammillaria parkinsonii* Ehrenb. (Cactaceae)
Morphoanatomy, histochemistry and germination of the seeds of Mammillaria parkinsonii Ehrenb. (Cactaceae)
Y. Uribe Salazar | A. Quintanar Isaias | C. Barbosa Martínez | J. Flores | C.L. Jiménez Sierra
- 135 Asymbiotic germination, *ex situ* conservation and *in vitro* plant regeneration of *Catasetum integerrimum* Hook
Germinación asimbiótica, conservación ex situ e in vitro regeneración de plantas de Catasetum integerrimum Hook
G. López Puc | G.J. Herrera Cool
- 151 Fitoquímicos y propiedades nutraceuticas de durazno (*Prunus persica* L.) cultivado en Zacatecas
*Phytochemicals and nutraceutical properties of peach (*Prunus persica* L.) harvested in Zacatecas*
J. Aguayo Rojas | S. Mora Rochín | X. Tovar Jiménez | J.J. Rochín Medina | R.O. Navarro Cortez
- 167 Evaluation of extracts of endemic trees (*Magnolia* spp.) in Mexico against the fruit fly pest and preliminary phytochemical study
*Evaluación de extractos de árboles endémicos (*Magnolia* spp.) de México contra la plaga de la mosca de la fruta y estudio fitoquímico preliminar*
S.G. Vásquez Morales | E.A. Alvarez Vega | D.A. Infante Rodríguez | J.P. Huchin Mian | M. Pedraza Reyes
- 183 Características fenotípicas, nutricionales y nutraceuticas de frutos de chile x'catik, dulce y su híbrido f1 (*Capsicum annuum* L.)
*Phenotypic, nutritional and nutraceutical traits of x'catik chili fruits, sweet and its f1 hybrid (*Capsicum annuum* L.)*
Y.A. Mis Valdez | M.J. Hernández Pinto | R. Garruña | K.B. Medina Dzul | R.H. Andueza Noh
- 197 Mecanismos de infección endógena en frutos de cacao con *Moniliophthora roreri*
Mechanisms of endogenous infection in cocoa fruits with Moniliophthora roreri
V. Flores | L. Gómez Rodríguez | J.A. López García | J. Grajales Conesa
- 211 Efectos de *Bacillus subtilis* cepas GBO3 y IN937b en el crecimiento de maíz (*Zea mays* L.)
*Effects of *Bacillus subtilis* strains GBO3 and IN937b on the growth of corn (*Zea mays* L.)*
A.E. Gutiérrez Calvo | A. Gutiérrez Estrada | C.L. Miceli Méndez | M.A. López Miceli
- 219 Conhecimentos etnobotânicos de mateiros em comunidades rurais da região da Serra das Almas, Paraná - Brasil
Conocimiento etnobotánico de los silvicultores en comunidades rurales de la región Serra das Almas, Paraná - Brasil
Ethnobotanical knowledge of foresters in rural communities in the Serra das Almas region, Paraná - Brazil
M. Ferreira Clarindo | A. Staniski | J. Strachulski
- 239 Valor cultural de los recursos forestales no maderables en comunidades zapotecas de la Sierra Juárez de Oaxaca
Cultural value of non-timber forest resources in Zapotec communities of the Sierra Juarez de Oaxaca
J. Martínez López | N.G. Molina Luna | S. Rangel Landa | C. Aquino Vázquez | A. Acosta Ramos



Portada

Podopterus mexicanus Humb. & Bonpl. Polygonaceae. “Rompe capa”. Árboles o arbustos de 1.5-6.0 m de altura, ramas con espina terminal, braquiblastos, hojas fasciculadas, flores blancas a verdosas en fascículos, y frutos cubiertos por el perianto externo que forma alas delgadas y largamente decurrentes hacia el pedicelo. Crece en bosques tropicales caducifolios y bosques espinosos, sobre suelo rocoso negro derivado de rocas ígneas. En elevaciones de 550-760 m. Florece de abril a mayo y fructifica de junio a septiembre. Se distribuye desde México hasta Centroamérica. En México se encuentra en los estados de Colima, Guerrero, Oaxaca, Puebla, Tamaulipas, Veracruz y Yucatán. Se utiliza como planta melífera y para leña.

Podopterus mexicanus Humb. & Bonpl. Polygonaceae. "Rompe capa". Trees or shrubs 1.5-6.0 m tall, branches with a terminal spine, brachyblasts, fasciculate leaves, white to greenish flowers in fascicles, and fruit covered by the external perianth that forms thin wings and longly decurrent towards the pedicel. It grows in tropical deciduous forests and thorny forests, on black, rocky soil derived from igneous rocks. At elevations of 550-760 m. Blossoms from April to May and bears fruit from June to September. It is distributed from Mexico to Central America. In Mexico it is found in the states of Colima, Guerrero, Oaxaca, Puebla, Tamaulipas, Veracruz and Yucatán. It is used as a melliferous plant and for firewood.

por/by Rafael Fernández Nava



INSTITUTO POLITÉCNICO NACIONAL

Director General: *Dr. Arturo Reyes Sandoval*

Secretario General: *Mtro. Juan Manuel Cantú Vázquez*

Secretario Académico: *Dr. David Jaramillo Vigueras*

Secretario de Extensión e Integración Social: *Dr. Luis Alfonso Villa Vargas*

Secretario de Investigación y Posgrado: *Dra. Laura Arreola Mendoza*

Secretario de Servicios Educativos: *Dra. Ana Lilia Coria Páez*

Secretario de Administración: *M. en C. Javier Tapia Santoyo*

Director de Educación Superior: *Mtro. Mauricio Igor Jasso Zaranda*

ESCUELA NACIONAL DE CIENCIAS BIOLÓGICAS

Directora:

Dra. Yadira Rivera Espinoza

Subdirectora Académica:

M. en C. Martha Patricia Cervantes Cervantes

Subdirector Administrativo:

Ing. Raúl Chávez Alvircio

Jefe de la Sección de Estudios de Posgrado e Investigación:

Dr. Gerardo Aparicio Ozores

Subdirector de Servicios Educativos e Integración Social:

Dr. Felipe Neri Rodríguez Casasola

POLIBOTÁNICA, Año 27, No. 53, enero-junio 2022, es una publicación semestral editada por el Instituto Politécnico Nacional, a través de la Escuela Nacional de Ciencias Biológicas. Unidad Profesional Lázaro Cárdenas, Prolongación de Carpio y Plan de Ayala s/n, Col. Santo Tomás C.P. 11340 Delegación Miguel Hidalgo México, D.F. Teléfono 57296000 ext. 62331.

<http://www.herbario.encb.ipn.mx/>, Editor responsable: Rafael Fernández Nava. Reserva de Derechos al Uso Exclusivo del Título No. 04-2015-011309001300-203. ISSN impreso: 1405-2768, ISSN digital: 2395-9525, ambos otorgados por el Instituto Nacional del Derecho de Autor. Responsable de la última actualización de este número, Unidad de informática de la ENCB del IPN, Rafael Fernández Nava, Unidad Profesional Lázaro Cárdenas, Prolongación de Carpio y Plan de Ayala s/n, Col. Santo Tomás C.P. 11340 Delegación Miguel Hidalgo México, D.F.

Las opiniones expresadas por los autores no necesariamente reflejan la postura del editor de la publicación.

Queda estrictamente prohibida la reproducción total o parcial de los contenidos e imágenes de la publicación sin previa autorización del Instituto Politécnico Nacional.

REVISTA BOTÁNICA INTERNACIONAL DEL INSTITUTO POLITÉCNICO NACIONAL

EDITOR EN JEFE
Rafael Fernández Nava

EDITORAS ASOCIADAS
Maria de la Luz Arreguín Sánchez

COMITÉ EDITORIAL INTERNACIONAL

Christiane Anderson
University of Michigan
Ann Arbor. Michigan, US

Edith V. Gómez Sosa
Instituto de Botánica Darwinion
Buenos Aires, Argentina

Heike Vibrans
Colegio de Postgraduados
Estado de México, México

Jorge Llorente Bousquets
Universidad Nacional Autónoma de México
Ciudad de México, México

Graciela Calderón de Rzedowski
Instituto de Ecología del Bajío
Pátzcuaro, Mich., México

Delia Fernández González
Universidad de León
León, España

Theodore S. Cochran
University of Wisconsin
Madison, Wisconsin, US

Jerzy Rzedowski Rotter
Instituto de Ecología del Bajío
Pátzcuaro, Mich., México

Hugo Cota Sánchez
University of Saskatchewan
Saskatoon, Saskatchewan, Canada

Luis Gerardo Zepeda Vallejo
Instituto Politécnico Nacional
Ciudad de México, México

Fernando Chiang Cabrera
Universidad Nacional Autónoma de México
Ciudad de México, México

Claude Sastre
Muséum National d'Histoire Naturelle
París, Francia

Thomas F. Daniel
California Academy of Sciences
San Francisco, California, US

Mauricio Velayos Rodríguez
Real Jardín Botánico
Madrid, España

Francisco de Asís Dos Santos
Universidad Estadual de Feira de Santana
Feira de Santana, Brasil

Noemí Waksman de Torres
Universidad Autónoma de Nuevo León
Monterrey, NL, México

Carlos Fabián Vargas Mendoza
Instituto Politécnico Nacional
Ciudad de México, México

Julieta Carranza Velázquez
Universidad de Costa Rica
San Pedro, Costa Rica

José Luis Godínez Ortega
Universidad Nacional Autónoma de México
Ciudad de México, México

Tom Wendt
University of Texas
Austin, Texas, US

José Manuel Rico Ordaz
Universidad de Oviedo
Oviedo, España

DISEÑO Y FORMACIÓN ELECTRÓNICA
Luz Elena Tejeda Hernández

OPEN JOURNAL SYSTEM Y TECNOLOGÍAS DE LA INFORMACIÓN
Pedro Aráoz Palomino

Toda correspondencia relacionada con la revista deberá ser dirigida a:

Dr. Rafael Fernández Nava

Editor en Jefe de

POLIBOTÁNICA

Departamento de Botánica

Escuela Nacional de Ciencias Biológicas, Instituto Politécnico Nacional
Apdo. Postal 17-564, CP 11410, Ciudad de México

Correo electrónico:

polibotanica@gmail.com

rfernán@ipn.mx

Dirección Web

http://www.polibotanica.mx

POLIBOTÁNICA es una revista indexada en:

CONACYT, índice de Revistas Mexicanas de Investigación Científica y Tecnológica del Consejo Nacional de Ciencia y Tecnología.

SciELO - Scientific Electronic Library Online.

Google Académico - Google Scholar.

DOAJ, Directorio de Revistas de Acceso Público.

Dialnet portal de difusión de la producción científica hispana.

REDIB Red Iberoamericana de Innovación y Conocimiento Científico.

LATINDEX, Sistema regional de información en línea para revistas científicas de América Latina, el Caribe, España y Portugal.

PERIODICA, Indice de Revistas Latinoamericanas en Ciencias.



Polibotánica

ISSN electrónico: 2395-9525

polibotanica@gmail.com

Instituto Politécnico Nacional

México

<http://www.polibotanica.mx>

**ASYMBIOTIC GERMINATION,
in vitro CONSERVATION AND
REGENERATION OF *Catasetum*
integerrimum.**

**GERMINACIÓN ASIMBIÓTICA,
CONSERVACIÓN Y
REGENERACIÓN *in vitro* DE
Catasetum integerrimum.**

López Puc G. y Herrera-Cool G. J.

ASYMBIOTIC GERMINATION, *in vitro* CONSERVATION AND REGENERATION OF *Catasetum integerrimum*.

GERMINACIÓN ASIMBIÓTICA, CONSERVACIÓN Y REGENERACIÓN *in vitro* DE *Catasetum integerrimum*.



**ASYMBIOTIC GERMINATION, *in vitro* CONSERVATION AND REGENERATION OF
*Catasetum integerrimum***

**GERMINACIÓN ASIMBIÓTICA, CONSERVACIÓN Y REGENERACIÓN *in vitro*
DE *Catasetum integerrimum***

López Puc G.
y Herrera-Cool G. J.

ASYMBIOTIC
GERMINATION, *in vitro*
CONSERVATION AND
REGENERATION OF
Catasetum integerrimum.

GERMINACIÓN
ASIMBIÓTICA,
CONSERVACIÓN Y
REGENERACIÓN *in vitro*
DE *Catasetum integerrimum*.

POLIBOTÁNICA
Instituto Politécnico Nacional

Núm. 53: 135-149. Enero 2022

DOI:
10.18387/polibotanica.53.9

G. López Puc / glopez@ciatej.mx

*Centro de Investigación y Asistencia en Tecnología y Diseño del Estado de Jalisco,
Subsede Sureste, Sierra Papacal, Yucatán, 97302, México.*

G.J. Herrera-Cool

*Campo Experimental Chetumal, INIFAP,
Chetumal-Bacalar Km 25, 77900, México.*

RESUMEN: La deforestación y recolecta ilegal de las orquídeas ha generado la necesidad de plantear alternativas para conservarlas y multiplicarlas. Una orquídea de interés es *Catasetum integerrimum*, cuyo hábitat está en América Central y México. El objetivo de este estudio fue proporcionar métodos para la conservación y uso sustentable de *C. integerrimum*. Se establecieron las condiciones para la germinación asimbiótica de semillas mediante la evaluación de cuatro formulaciones de medios basales: 1) Murashige y Skoog (MS) con macro y micronutrientes y vitaminas; 2) MS con macro y micronutrientes sin vitaminas; 3) Medio basal para orquídeas con carbón activado y polvo de plátano y 4) Medio basal Knudson. Para la conservación *in vitro* del germoplasma se desarrolló un protocolo de crecimiento lento para lo cual se evaluó el efecto de la concentración del medio basal MS (Fuerza iónica completa y a la mitad de su fuerza iónica) y el efecto del sorbitol, manitol y sacarosa a 1, 2 y 3%. Con el fin de establecer un protocolo para la regeneración *in vitro* para la multiplicación de plantas se evaluó el efecto de tres tipos de explante (raíz, nudo, y hoja) y el efecto individual de N6-furfuriladenina (kinetina) a 4.64, 9.29 y 23.2 µM y 6- Bencil Amino Purina (BAP) a 4.43, 8.87 y 22.2 µM. Los resultados mostraron que la germinación asimbiótica de *C. integerrimum* fue exitosa en el medio Murashige y Skoog con macro y micronutrientes sin vitaminas. La conservación *in vitro* fue posible en el medio basal MS a la mitad de su fuerza iónica suplementado con sorbitol al 2%, ya que en este tratamiento las plantas mostraron menor crecimiento después de 6 meses de cultivo. La organogénesis directa de brotes se observó en el explante de nudo en MS suplementado con BAP y en MS con kinetina en todas las concentraciones evaluadas. La organogénesis indirecta de brotes se observó en explante de raíz en MS suplementado con 4.64 o 9.29 µM de kinetina.

Palabras clave: lento crecimiento; kinetina; manitol; sorbitol, conservación a mediano plazo.

ABSTRACT: The deforestation and illegal over-collection of orchids have generated the need to propose alternatives to conserve and multiply them. One orchid of interest is *Catasetum integerrimum*, whose habitat is in Central America and Mexico. This study aimed to provide methods for the conservation and sustainable use of *C. integerrimum*. Asymbiotic seed germination was established by evaluating four formulations of basal media: 1) Murashige and Skoog (MS) with macro and micronutrients and vitamins; 2) MS with macro and micronutrients without vitamins; 3) Basal medium for orchids with activated carbon and banana powder; 4) Knudson basal medium. We developed the slow-growth protocol for which evaluated MS basal medium concentration (full and half of the ionic strength) and the effect of sorbitol, mannitol, and sucrose at 1, 2, and 3%. The influence of three explant types (root, node,

and leaf) and kinetin at 4.64, 9.29, and 23.2 μM and Benzyl Amino Purine (BAP) at 4.43, 8.87, and 22.2 μM were evaluated to establish *in vitro* regeneration for plant multiplication. Asymbiotic germination of *C. integerrimum* was successful on Murashige and Skoog medium with macro and micronutrients without vitamins. *In vitro* conservation was viable in half-strength basal MS medium supplemented with 2% sorbitol; it was the best treatment for conserving seedlings under slow-growth conditions after six months of *in vitro* culture. Node explant formed direct shoot organogenesis in MS with BAP or kinetin at all concentrations evaluated. Indirect shoot organogenesis was observed in root explant in MS supplemented with 4.64 or 9.29 μM kinetin.

Keywords: slow growth; kinetin, mannitol; sorbitol, medium-term conservation

INTRODUCTION

Orchidaceae is one of the largest families of flowering plants (Pérez Gutiérrez, 2010; Noguera-Savelli & Cetzel-Ix, 2014), which are part of complex biological interactions of ecosystems (Seaton *et al.*, 2013; Pereira *et al.*, 2015). Orchids are also considered an important crop in the floriculture industry. This plant family is usually cultured due to its medicinal properties and uses in traditional food, as well as the generation of nutraceutical products, perfumes, and cosmetics (Hinsley *et al.*, 2018; Kanlayavattanakul *et al.*, 2018; Salazar-Mercado & Botello-Delgado, 2020; Singh *et al.*, 2021). In recent years, many species of orchids have been threatened in their natural habitats because of the environmental pressure generated by humans (Juras *et al.*, 2020; Fonseka, 2020; Puspitaningtyas & Handini, 2020). *C. integerrimum* is an orchid that reaches to adult stage in approximately two years. This development stage is less than 50 percent of the time required by most of the not deciduous taxa orchids (Benzing *et al.*, 1982). This species lives in Guatemala, Belize, Honduras, El Salvador, Nicaragua and Mexico (Morales-Báez *et al.*, 2016). *C. integerrimum* is an ornamental plant whose flowers are very showy and attractive that produce a fragrance compound by pinene (32.52%) and carvone oxide (46.21%) (del Mazo Cancino & Damon, 2007). It is often collected, resulting in a decrease in the population of their natural habitat (Morales-Báez *et al.*, 2016). This over-exploitation generates a loss of their population. Moreover, germination in a natural environment entirely depends on the mycorrhizal fungi association (Pérez Gutiérrez, 2010). This association provides water, carbohydrates, minerals, and vitamins for embryos development; however, mortality rates of embryos developed are high despite this association (Kauth *et al.*, 2006). Asymbiotic seed propagation, which is an excellent method for the micropropagation of orchids for conservation purposes (Kim *et al.*, 2021). Asymbiotic germination of orchid seeds was established by Lewis Knudson (1884–1958), and it was the first practical procedure for *in vitro* propagation of any plant in pure (i.e., axenic) culture (Yam & Arditti, 2009). Since then, many ornamental species of orchids have been micropropagated using this method (Pereira *et al.*, 2017; Vudala & Ribas, 2017; Acemi & Özen, 2019; Franceschi *et al.*, 2019; Koene *et al.*, 2019; Fonseka, 2020; Kang *et al.*, 2020; Salazar-Mercado & Botello-Delgado, 2020; Sorgato *et al.*, 2020; Chin *et al.*, 2021; Kim *et al.*, 2021; Navarro *et al.*, 2021; Yeh *et al.*, 2021); The asymbiotic germination process generates protocorm-like-bodies (PLBs) which are embryogenic structures (Chin *et al.*, 2021). The process goes from stage 0: No germination to stage 5: the differentiation (Yamazaki & Miyoshi, 2006).

Due to the conservation of the germplasm of this species is needed, *in vitro* culture techniques have been successfully applied for conservation. Also, it facilitates the availability of germplasm for use in genetic improvement programs (Arrigoni-Blank *et al.*, 2014).

One of the methods used is slow growth, which reduces plant metabolism and consequently decelerates its development, which allows the maintenance of *in vitro* plant collections without the need for subculture (Chauhan *et al.*, 2019; Santos Lima *et al.*, 2021; Petrova *et al.*, 2021). The slow growth procedure is called medium-term conservation or minimal growth storage (Muñoz *et al.*, 2019).

The advantage of the slow growth technique is that enabling *in vitro* maintenance of shoot cultures under aseptic conditions by markedly reducing the frequency of periodic subculturing without affecting the viability and regrowth potential of such shoot cultures (Koeda *et al.*, 2018). The slow growth can be achieved by basal culture medium modification, sucrose, and plant growth regulators (PGRs) concentration changes (Ruta *et al.*, 2020). Other substances used to slow growth are mannitol and sorbitol (da Costa Santos *et al.*, 2011; Tavazza *et al.*, 2015; Hammond Hammond *et al.*, 2019; Muñoz *et al.*, 2019; Petrova *et al.*, 2021;). Furthermore, reports of slow-growth techniques include protocols for conserved material multiplication (Hammond-Hammond *et al.*, 2019; Ramírez-Mosqueda *et al.*, 2019; Fonseka, 2020).

The present study describes the asymbiotic germination of *C. integerrimum*, its conservation in a slow-growth medium and the regeneration by direct and indirect adventitious organogenesis.

MATERIAL AND METHODS

Obtaining plant material

Fully closed of *C. integerrimum* from two capsules with good condition were collected from experimental field El Tormento, Campeche, Mexico at the "Instituto Nacional de Investigaciones Forestales Agrícolas y Pecuarias (INIFAP)". Afterward, the capsules collected were transferred to the laboratory and put inside a desiccator with 1 kg of silica gel at $25\pm2^{\circ}\text{C}$ for five days. Then they were washed with a solution of Extran® at 5% for 5 minutes and rinsed three times with sterile distillate water.

In vitro establishment of plant material

The capsules were disinfected in a laminar flow cabinet by immersion in a solution of sodium hypochlorite at 1% with 10 µL of Extran® for 10 min, then were rinsed three times with distilled water, afterward were atomized with 70% of ethanol and flamed with a burner. The capsules sterilized were longitudinally cut with a scalpel to obtain the seeds, which were put on a sterile Petri dish. Finally, 300 seeds were cultured in Magenta® boxes with treatments for asymbiotic germination.

Asymbiotic germination

Semi-solid media of MS medium with macro-micro nutrients and vitamins (M5519 SIGMA®), MS medium with macro-micro nutrients (M5524 SIGMA®), Orchid Medium with charcoal and Banana Powder (P1056 SIGMA®), and Knudson C (K4003 SIGMA®) were used to investigate the asymbiotic germination of *C. integerrimum* without PGRs. The pH of the media was adjusted at 5.8 before sterilizing in an autoclave at 121°C . The treatments were dosed at 25 mL per Magenta® box, and seeds were established in three Magenta® boxes per treatment. The evaluation was carried out every 4 weeks for 16 weeks, recording data and digital pictures.

Slow growth for *in vitro* conservation

Plantlets of 2 cm from asymbiotic germination were established in a factorial experiment of 2 x 3 x 3 for conservation. The first factor was the concentration of basal medium (Full and half ionic strength), the second factor was the carbon source (sorbitol, mannitol, and sucrose), and the third factor was the concentration of carbon source (1, 2, and 3%). The experiment was carried out with two MS basal medium control: half-strength MS basal medium (2.2 g L^{-1}) and full-strength MS basal medium (4.4 g L^{-1}) without carbon source. In total were evaluated 18 treatments with two controls and three replicates per treatment. Treatments with half-strength MS basal medium were solidified with 3.1 g L^{-1} of Gelrite® and the treatments with full strength MS basal medium with 2.2 g L^{-1} of Gelrite®. The variables evaluated were the plantlet growth and the shoots number formed per treatment. Plant growth was obtained by subtracting the plantlet length (mm) at six months of culture from plantlet length at the beginning of

culture. The result was the plantlet growing after six months of culture. The shoots number formed per treatment was recorded at 12 months.

Plant regeneration

Induction of regeneration was evaluated in a factorial design of $3 \times 2 \times 3$, which the first factor was the explant source (node, leaf, and root), the second factor was kinetin or BAP, and the third factor PGR concentration, kinetin at 4.64, 9.29 and 23.2 μM , and BAP at 4.43, 8.87 and 22.2 μM . The experiment had 18 treatments and one control with three replicates per treatment. The control was the MS basal medium without PGRs. All treatments were formulated with full strength MS basal medium added with 3% sucrose and 2.2g L^{-1} of Gelrite®. The variable evaluated was shoot number per explant at the 60 days, and shoots were allowed to grow until 120 days of culture. All treatments were supplemented with 2 g L^{-1} of activated carbon.

In vitro culture conditions

All treatments were incubated at $23 \pm 2^\circ\text{C}$ and relative humidity of 60% with a photoperiod of 16/8 hours (light/darkness); the light source was provided by LED lamps with a light intensity of $60 \mu\text{mol}^2\text{s}^{-1}$.

Statistical analysis

Data were subjected to one-way ANOVA analysis of variance for comparison of means. Significant differences were calculated according to the Tukey test at the 5% significance level using a statistical Statgraphics® Centurion XVI statistical software. Data were presented as means \pm standard error.

RESULTS

Asymbiotic germination

Asymbiotic germination of seeds of *C. integerrimum* occurred in basal MS medium with macro and micronutrients without vitamins, 250 plantlets were obtained. At four weeks after sowing, the seeds formed a brown friable callous mass jelly-like, from which white embryogenic structures emerged (WES) (fig. 1 A). At six weeks, the WES began to grow into PLBs that turned green. (fig. 1 B). Root formation in PLBs (fig. 1 C) and shoot differentiation into PLBs from protocorms (fig. 1 D) occurred at 8 and 10 weeks, respectively. The plantlets formed rootlets and leaves at 12 weeks (fig. 1 E). The same basal medium where germination occurred completed growth up to 16 weeks and reseeding was unnecessary during this process (fig. 1 F). *C. integerrimum* seeds sown on basal MS media with macro and micronutrients and vitamins, Orchid Medium with Charcoal and Banana Powder and Knudson had no germination response.

Slow growth for *in vitro* conservation

There was significant difference among the 20 treatments selected to evaluate the growth of *C. integerrimum* plantlets (Table 1). T3 allows the highest growth value (76.24 ± 30.5) followed by T1 (54.74 ± 7.6), T5 (52.69 ± 20.1), T4 (48.32 ± 15.8), and T2 (37.39 ± 3.7) (Table 1, fig. 1). The treatments with the lowest growth values of cultivated plantlets were T19 (8.54 ± 5.2), T20 (13.10 ± 5.3), T17 (14.18 ± 6.1), T18 (17.70 ± 5.8), T16 (18.61 ± 4.0), T10 (19.71 ± 2.6) and T11 (21.07 ± 8.6), there was no significant difference among these seven treatments (Table 1), the controls T19 and T20 showed the lowest growth values, the leaves of the plantlets showed etiolation and died. Treatment T17 was selected because it presented the lowest value of growth in plantlet length among the other treatments. The plantlets from T17 did not show changes in plant morphology. At six months the shoot formation in the slow growth treatments was evaluated. The treatment T3 (Table 1) formed the highest number of shoots (28.4 ± 3.2). On the other hand, T17 showed the lowest shoot formation (1 ± 0.0).




Fig. 1. Asymbiotic seed germination of *C. integerrimum*. Callus with embryogenic structures (A); Embryogenic structures transforming into PLBs (B); Rhizoids formed on the protocorm (white arrows) (C); Shoots differentiated from the protocorm (D); Plantlets formed 3 months after sowing seeds (E); Fully developed plantlets at four months of age (F).

Table 1. Effect of culture medium composition on plantlets growth and shoot proliferation of *C. integerrimum* Hook. to determine the best treatment for *in vitro* conservation

Treatment	MS concentration (%)	Carbon source	Carbon concentration (%)	Average of plantlet length (mm)	No. Shoot per treatment
1	100	Sucrose	1	54.74±7.6 ^h	9.20±2.2 ^c
2			2	37.39±3.7 ^{f,g}	17.60±2.7 ^e
3			3	76.24±30.5 ^j	28.40±3.2 ^f
4	50	Sucrose	1	48.32±15.8 ^{gh}	12.60±1.9 ^{cd}
5			2	52.69±20.1 ^h	14.40±2.3 ^{de}
6			3	26.84±8.8 ^{def}	5.40±1.3 ^b
7	100	Mannitol	1	31.95±13.1 ^{ef}	3.40±1.1 ^{ab}
8			2	32.26±6.0 ^{ef}	2.60±0.8 ^{ab}
9			3	24.23±4.5 ^{bcd}	2.20±0.4 ^{ab}
10	50	Mannitol	1	19.71±2.6 ^{abcde}	2.40±0.5 ^{ab}
11			2	21.07±8.6 ^{abcde}	3.40±1.1 ^{ab}
12			3	31.57±7.4 ^{ef}	3.20±1.7 ^{ab}
13	100	Sorbitol	1	29.31±5.1 ^{def}	2.60±0.8 ^{ab}
14			2	30.14±3.9 ^{def}	2.80±1.3 ^{ab}
15			3	27.51±8.7 ^{def}	2.60±0.5 ^{ab}
16	50	Sorbitol	1	18.61±4.0 ^{abcde}	2.60±1.8 ^{ab}
17			2	14.18±6.1 ^{abc}	1.00±0.0 ^{ab}
18			3	17.70±5.8 ^{abcd}	3.20±1.3 ^b
19	100	None	0	8.540±5.2 ^a	0.00±0.0 ^a
20			0	13.10±5.3 ^{ab}	0.00±0.0 ^a

Different letters indicate significant statistical differences according Tukey test ($P \leq 0.05$). MS: Murashige and Skoog basal medium.

Plant regeneration

The type of explant influenced shoot formation (Table 2). The leaf explant did not show regeneration. Shoot induction from node explant of *C. integerrimum* via direct organogenesis occurred at 30 days (fig. 3 A) on treatments T2-T7. There was no significant statistical difference among treatments; T5, T16, and T17 showed 3.0 adventitious shoots per explant, followed by T3 with 2.5 shoots (Table 2). Rooting of shoots was observed in the same induction medium; the plantlets morphology at 60 and 120 days was normal (fig. 3 B-C).

The root explant formed callus after 30 days of cultivation in full strength MS basal medium supplemented with kinetin at 4.64 or 9.29 µM (fig. 4 A). Indirect shoot organogenesis at 60 days of culture (fig. 4 B), the rooting of shoots occurred in the same induction medium. Plantlets showed an average height of 8 cm after 120 days of cultivation. (fig. 4 C). BAP did not induce organogenesis in root explant (Table 2).

Table 2. Effect of PGR and type of explant on the induction of adventitious organogenesis of *C. integerrimum* Hook.

Treatment	Explant	PGR	PGR Concentration (µM)	# adventitious shoots/explant
1	Node	None	0.00	0.00±0.00 ^a
2		Kinetin	4.64	2.25±1.20 ^b
3			9.29	2.50±1.20 ^b
4			23.2	2.25±1.50 ^b
5		BAP	4.43	3.00±0.81 ^b
6			8.87	2.00±1.15 ^b
7			22.2	1.75±0.50 ^b
8	Leaf	None	0.00	0.00±0.00 ^a
9		Kinetin	4.64	0.00±0.00 ^a
10			9.29	0.00±0.00 ^a
11			23.2	0.00±0.00 ^a
12		BAP	4.43	0.00±0.00 ^a
13			8.87	0.00±0.00 ^a
14			22.2	0.00±0.00 ^a
15	Root	None	0.00	0.00±0.00 ^a
16		Kinetin	4.64	3.00±0.81 ^b
17			9.29	3.00±0.00 ^b
18			23.2	0.00±0.00 ^a
19		BAP	4.43	0.00±0.00 ^a
20			8.87	0.00±0.00 ^a
21			22.2	0.00±0.00 ^a

Different letters indicate significant statistical differences according Tukey test ($P \leq 0.05$). PGR: Plant Growth Regulator, BAP: Benzyl Amino Purine. MS: Murashige and Skoog basal medium.




Fig. 2. *In vitro* conservation of *C. integerrimum* growth after 6 months of cultivation in treatments evaluated for its conservation. Full strength MS basal medium supplemented with sucrose 1, 2, and 3%: (1), (2), and (3) respectively. Half strength MS basal medium with sucrose 1, 2, and 3%: (4), (5), and (6) respectively. Full strength MS basal medium supplemented with mannitol 1, 2, and 3%: (7), (8), and (9) respectively. Half strength MS basal medium with mannitol 1, 2, and 3%: (10), (11), and (12) respectively. Full strength MS basal medium supplemented with sorbitol 1, 2, and 3%: (13), (14), and (15) respectively. Half strength MS basal medium supplemented with sorbitol 1, 2, and 3%: (16), (17), and (18) respectively. Controls: half-strength MS basal medium without carbon source (19) and full-strength MS basal medium without carbon source (20).




Fig. 3. Direct adventitious organogenesis from node explant of *C. integerrimum*. Adventitious shoots from node explant on full strength MS basal medium+ BAP at 4.43 μ M, 30 days culture (A); Plantlets at 60 days of culture (B); Plantlets at 120 days of culture (C).




Fig. 4. Indirect adventitious organogenesis from root explant of *C. integerrimum*. Callus formed in root explant at 30 days of culture (A); Adventitious shoots formed on the callus at 60 days of culture (B); Plantlets at 120 days of culture (C).

DISCUSSION

Tissue culture methods are an alternative for help in the rescue of species endangered by deforestation and unmeasured extraction from their natural habitat. This study presents asymbiotic germination, conservation, and *in vitro* regeneration of *C. integerrimum* that have not been previously reported.

The germination of orchid seeds in their natural habitat is complex and inefficient because there must be a symbiotic relationship with a fungus. Association between orchids and fungi was observed in 1824 (Yam & Arditti, 2009). On the other hand, *in vitro* germination is efficient when the composition of the culture medium is known.

According to the classification described by (Yam & Arditti, 2009), the stages of "pre-germination," "germination," and "protocorms formation" corresponding to stages 1-3 were not visible in this study. This observation could be due to the seeds cultivated on MS basal medium with macro and micronutrients without vitamins formed callus. Furthermore, white embryogenic structures formed from the callus, becoming PLBs, the stages 4 and 5 were clearly

visible. Full strength MS basal medium including all macro and micronutrients, without vitamins, was the only one of the four media evaluated that gave a response to asymbiotic germination of *C. integerrimum*. Unlike reports of asymbiotic germination using MS medium with half of the macro and micronutrients (Koene *et al.*, 2019; Kang *et al.*, 2020; Yeh *et al.*, 2021; Navarro *et al.*, 2021; Kim *et al.*, 2021). In vitro culture conditions and the lack of vitamins in the MS basal medium may have generated stress in the *C. integerrimum* seed culture, triggering a signaling cascade for dedifferentiation leading to callus formation. It is going by concepts reported by Fehér (2014) that differentiated plant cells, under certain circumstances, can revert to an earlier developmental state and regain totipotency.

Mineral nutrition, vitamins, and medium composition influence asymbiotic seed germination have been reported (Salazar-Mercado & Botello-Delgado, 2020; Kim *et al.*, 2021). Other formulations have been necessary, for instance: New Dogashima medium was used for germination of mature seeds of *Calanthe tricarinata* (Godo *et al.*, 2010) or Woody plant medium (WPM) (Vudala & Ribas, 2017; Franceschi *et al.*, 2019); Vacin and Went medium (VW) (Pereira *et al.*, 2017; Utami & Hariyanto, 2019; Acemi & Özen, 2019). New Dogashima medium has microelements and organic compounds of Nitsch and Morel respectively, NH_4NO_3 and KNO_3 are in minor amount in comparison with MS, this media was developed for micropropagation of orchids (Tokuhara *et al.*, 1993). On the other hand, Woody Plant Medium contains the macro-micronutrients and vitamins described by Phillips & Garda (2019).

In this study *C. integerrimum* exogen PGRs were not necessary during the asymbiotic germination and plantlet development contrary to other protocols reported, asymbiotic germination of *Calanthe tricarinata* Lindl occurred using the basal medium New Dogashima with BAP or α -naphthalene acetic acid (NAA) (Godo *et al.*, 2010); On the other hand, the germination of *C. integerrimum* was achieved using full-strength MS basal medium added with 2% BAP; in the same study the protocorms development was necessary to use full strength MS basal medium added with 5 mg L⁻¹ of activated charcoal and indole acetic acid (IAA) at 5 mg L⁻¹ (Velázquez Kú *et al.*, 2016).

The conservation protocols for *Dendrobium draconis*, *Bulbophyllum auricomum* Lindl., *Ullucus tuberosus* species of the subfamily Epidendroideae, *Arnica montana* L. used MS reduced to half its ionic strength as basal medium (Than, 2013; Hammond Hammond *et al.*, 2019; Menezes-Sá *et al.*, 2019; Petrova *et al.*, 2021).

Other basal media used in preservation protocols include Vacin and Went (Puspitaningtyas & Handini, 2020). The woody plant medium (WPM) (Matos Alvim *et al.*, 2020). Culture media for slow growth can use sucrose as a carbon source (Than, 2013; Sanghamitra *et al.*, 2019; Menezes-Sá *et al.*, 2019; El-Hawaz *et al.*, 2019; Matos Alvim *et al.*, 2020; Gomes *et al.*, 2021). Another modification made to the culture medium is sorbitol and mannitol as carbon sources. The use of mannitol on slow growth protocols has been reported for Globe artichoke, *Ullucus tuberosus* (Tavazza *et al.*, 2015; Hammond Hammond *et al.*, 2019). On the other hand, the combined effect of mannitol and sucrose allowed the conservation of *Amburana cearensis* plants with better characteristics than using sucrose alone (Matos Alvim *et al.*, 2020). The results showed that *C. integerrimum* plantlets have higher growth in treatments T1-T5 supplemented with sucrose, even inducing multiple sprouting, so they are recommended for propagation but not for slow growth. T17 treatment formulated with half-strength MS basal medium and 2% sorbitol was selected for *in vitro* conservation of *C. integerrimum*. The use of sorbitol to slow growth rate has been reported in *Hancornia speciosa* Gomes; *Arnica montana* L., *Solanum tuberosum* (da Costa Santos *et al.*, 2011; Muñoz *et al.*, 2019; Petrova *et al.*, 2021); The combination of sorbitol with sucrose preserves *in vitro* plants of *Solanum tuberosum* without phenotypic abnormalities (Nasiruddin & Rafiul Islam, 2018). The combination of sorbitol with sucrose and mannitol with sucrose allows slow growth storage of potato germplasm (Muñoz *et al.*, 2019).

In this study, kinetin induced the regeneration of adventitious shoots from root explants, and BAP induced a similar response from node explants. Kinetin induces callus (clusters of dedifferentiated plant cells) to differentiate into adventitious buds (Barciszewski *et al.*, 2007; Sianipar & Mariska, 2020; Sharma *et al.*, 2021).

CONCLUSIONS

Asymbiotic germination, *in vitro* conservation, and adventitious shoots regeneration of *C. integerrimum* are methods effective and efficient. It is also important to highlight that the *in vitro* conservation protocol has been so effective that it has maintained this species *in vitro* conservation for ten years. The asymbiotic germination method and the regeneration method can be used for propagation, helping to reduce the collection of this species in its habitat.

LITERATURE CITED

- Acemi, A., & Özen, F. (2019). Optimization of *in vitro* asymbiotic seed germination protocol for *Serapias vomeracea*. *The EuroBiotech Journal*, 3(3), 143–151. <https://doi.org/10.2478/ebtj-2019-0017>
- Arrigoni-Blank, M. D. F., Tavares, F. F., Blank, A. F., Santos, M. C. dos, Menezes, T. S. A., & Santana, A. D. D. de. (2014). *In vitro* conservation of sweet potato genotypes. *The Scientific World Journal*, 2014, 1–7. <https://doi.org/10.1155/2014/208506>
- Barciszewski, J., Massino, F., & Clark, B. F. C. (2007). Kinetin-A multiactive molecule. *International Journal of Biological Macromolecules*, 40(3), 182–192. <https://doi.org/10.1016/j.ijbiomac.2006.06.024>
- Benzing, D. H., Bent, A., Moscow, D., Peterson, G., & Renfrow, A. (1982). Functional correlates of deciduousness in *Catasetum integerrimum* (Orchidaceae). *Selbyana*, 7(1), 1–9.
- Chin, C. K., Stanly, C., Chew, B. L., & Subramaniam, S. (2021). Modified basal culture medium improves proliferation of *Dendrobium Sabin Blue's* protocorm-like bodies (PLBs). *Biologia*, 76, 1433–1443. <https://doi.org/10.1007/s11756-021-00743-8/Published>
- da Costa Santos, M., da Silva Lédo, A., da Silva Lédo, C. A., Duarte Souza, F. V., & da Silva Junior, J. F. (2011). Effect of sucrose and sorbitol in the *in vitro* conservation of mangaba tree nodal segments. *Revista Ciência Agronômica*, 42(3), 735–741.
- del Mazo Cancino, A., & Damon, A. (2007). Fragrance analysis of euglossine bee pollinated orchids from Soconusco, south-east Mexico. *Plant Species Biology*, 22(2), 129–134. <https://doi.org/10.1111/j.1442-1984.2007.00185.x>
- El-Hawaz, R. F., Adelberg, J., Naylor-Adelberg, J., Eisenreich, R., & van der Meij, J. (2019). The effect of slow-growth strategy on a production of *Petunia × hybrida* Vilm. microcuttings. *In vitro Cellular and Developmental Biology - Plant*, 55(4), 433–441. <https://doi.org/10.1007/s11627-019-09996-0>
- Fehér, A. (2014). *Somatic embryogenesis-Stress-induced remodeling of plant cell fate*. <https://doi.org/10.1016/j.bbagr.2014.07.005>
- Fonseka, K. (2020). Asymbiotic Seed Germination, Mass propagation and Conservation of Fox-tail orchid, *Rhynchostylis retusa* L. Blume: An endangered orchid. *Asian Journal of Conservation Biology*, 9(2), 144.
- Franceschi, C. R. B., Smidt, E. C., Vieira, L. N., & Ribas, L. L. F. (2019). Storage and *in vitro* germination of orchids (Orchidaceae) seeds from Atlantic forest – Brazil. *Anais Da Academia Brasileira de Ciencias*, 91(3). <https://doi.org/10.1590/0001-3765201920180439>
- Godo, T., Komori, M., Nakaoki, E., Yukawa, T., & Miyoshi, K. (2010). Germination of mature seeds of *Calanthe tricarinata* Lindl., an endangered terrestrial orchid, by asymbiotic

- culture *in vitro*. *In vitro Cellular and Developmental Biology - Plant*, 46(3), 323–328. <https://doi.org/10.1007/s11627-009-9271-1>
- Gomes, F., Clemente, M., Figueiredo, P., & Costa, R. L. (2021). Medium-term *in vitro* conservation of *Castanea* spp. hybrid clones. *Vegetos*, 34(1), 127–137. <https://doi.org/10.1007/s42535-020-00184-9>
- Hammond, S. D., Viehmannova, I., Zamecnik, J., Panis, B., & Hlasna Cepkova, P. (2019). Efficient slow-growth conservation and assessment of clonal fidelity of *Ullucus tuberosus* Caldas microshoots. *Plant Cell, Tissue and Organ Culture*, 138(3), 559–570. <https://doi.org/10.1007/s11240-019-01653-z>
- Hinsley, A., de Boer, H. J., Fay, M. F., Gale, S. W., Gardiner, L. M., Gunasekara, R. S., Kumar, P., Masters, S., Metusala, D., Roberts, D. L., Veldman, S., Wong, S., & Phelps, J. (2018). A review of the trade in orchids and its implications for conservation. In *Indonesia Botanical Journal of the Linnean Society* (Vol. 186). <https://academic.oup.com/botlinnean/article/186/4/435/4736317>
- Juras, M. C. R., Purgatto, E., Ferreira, W. de M., & Suzuki, R. M. (2020). Direct organogenesis and ethylene regulators in the cloning of *Epidendrum denticulatum* (Orchidaceae). *South African Journal of Botany*, 131, 374–379. <https://doi.org/10.1016/j.sajb.2020.03.010>
- Kang, H., Kang, K. W., Kim, D. H., & Sivanesan, I. (2020). *In vitro* Propagation of *Gastrochilus matsuran* (Makino) Schltr., an Endangered Epiphytic Orchid. *Plants*, 524, 1–10. <https://doi.org/10.3390/plants9040524>
- Kanlayavattanakul, M., Lourith, N., & Chaikul, P. (2018). Biological activity and phytochemical profiles of *Dendrobium*: A new source for specialty cosmetic materials. *Industrial Crops and Products*, 120, 61–70. <https://doi.org/10.1016/j.indcrop.2018.04.059>
- Kauth, P. J., Vendrame, W. A., & Kane, M. E. (2006). *In vitro* seed culture and seedling development of *Calopogon tuberosus*. *Plant Cell, Tissue and Organ Culture*, 85(1), 91–102. <https://doi.org/10.1007/s11240-005-9055-1>
- Kim, J. H., Son, S. W., Kim, S. Y., & Jeong, M. J. (2021). Asymbiotic seed germination and *in vitro* seedling development of *Pelatantheria scolopendrifolia*, a rare epiphytic orchid native to Korea. *Rhizosphere*, 19, 100371. <https://doi.org/10.1016/j.rhisph.2021.100371>
- Koeda, S., Matsumoto, S., Matsumoto, Y., Takisawa, R., Nishikawa, K., & Kataoka, K. (2018). Medium-term *in vitro* conservation of virus-free parthenocarpic tomato plants. *In vitro Cellular and Developmental Biology - Plant*, 54(4), 392–398. <https://doi.org/10.1007/s11627-018-9906-1>
- Koene, F. M., Amano, & Ribas, L. L. F. (2019). Asymbiotic seed germination and *in vitro* seedling development of *Acianthera prolifera* (Orchidaceae). *South African Journal of Botany*, 121, 83–91. <https://doi.org/10.1016/j.sajb.2018.07.019>
- Matos Alvim, B. F., Vieira de Souza, A. V., Lima-Brito, A., Tavares Fonseca, P., Soares, T. L., & Ferreira de Santana, J. R. (2020). *In vitro* conservation of *Amburana cearensis* (Fabaceae). *Ciencia Rural*, 50(7), 1–8. <https://doi.org/10.1590/0103-8478cr20190729>
- Menezes-Sá, T. S. A., Arrigoni-Blank, M. de F., da Costa, A. S., Feitosa-Alcantara, R. B., Blank, A. F., & Luz, J. M. Q. (2019). *In vitro* conservation and acclimatization of epidendroideae (Orchidaceae) from Sergipe, Brazil. *Bioscience Journal*, 35(2), 356–366. <https://doi.org/10.14393/BJ-v35n2a2019-38776>
- Morales-Báez, M., Salinas-Castro, A., Bello, D. E., Cadena, M. G. L., Fernández, A. R., & Trigos, A. (2016). *Stethobaroides nudiventris* (Coleoptera: Curculionidae), the Curculionid Cause of Petal Wilting on the *Catasetum integerrimum* Orchid. *Annals of the Entomological Society of America*, 109(6), 845–849. <https://doi.org/10.1093/aesa/saw057>
- Muñoz, M., Díaz, O., Reinún, W., Winkler, A., & Quevedo, R. (2019). Slow growth *in vitro* culture for conservation of chilotanum potato germplasm. *Chilean Journal of Agricultural Research*, 79(1), 26–35. <https://doi.org/10.4067/S0718-58392019000100026>

- Nasiruddin, M., & Rafiul Islam, A. (2018). *In vitro* slow growth conservation for two genotypes of *Solanum tuberosum* L. *Bangladesh J. Bot.*, 47(3), 369–380.
- Navarro, Q. R., de Oliveira Corrêa, D., Behling, A., Noseda, M. D., Amano, É., Suzuki, R. M., & Ribas, L. L. F. (2021). Efficient use of biomass and extract of the microalga *Desmodesmus subspicatus* (Scenedesmaceae) in asymbiotic seed germination and seedling development of the orchid *Cattleya warneri*. *Journal of Applied Phycology*. <https://doi.org/10.1007/s10811-021-02442-y>
- Noguera-Savelli, E., & Cetzel-Ix, W. (2014). Revisión e integración del conocimiento de las orchidaceae de Tabasco, México. *Botanical Sciences*, 92(4), 519–540.
- Pereira, G., Albornoz, V., Muñoz-Tapia, L., Romero, C., & Atala, C. (2015). Asymbiotic germination of *Bipinnula fimbriata* (Orchidaceae) seeds in different culture media. *Seed Science and Technology*, 43(3), 367–377. <https://doi.org/10.15258/sst.2015.43.3.01>
- Pereira, G., Albornoz, V., Romero, C., Lara, S., Sánchez-Olate, M., Ríos, D., & Atala, C. (2017). Asymbiotic germination in three *Chloraea* species (Orchidaceae) from Chile Germinación asimbiótica en tres especies de *Chloraea* (Orchidaceae) de Chile. *Gayana Bot.*, 74(1), 131–139.
- Pérez Gutiérrez, R. M. (2010). Orchids: A review of uses in traditional medicine, its phytochemistry and pharmacology. In *Journal of Medicinal Plants Research* 4(8), 592–638. <https://doi.org/10.5897/JMPR10.012>
- Petrova, M., Zayova, E., Geneva, M., Dimitrova, L., Vitkova, A., & Stanilova, M. (2021). Multiplication and Conservation of Threatened Medicinal Plant *Arnica montana* L. *Agric. Conspec. Sci.*, 86(1).
- Phillips, G. C., & Garda, M. (2019). Plant tissue culture media and practices: an overview. In *In vitro Cellular and Developmental Biology - Plant* 55(3), 242–257. Springer New York LLC. <https://doi.org/10.1007/s11627-019-09983-5>
- Puspitaningtyas, D. M., & Handini, E. (2020). *Ex-situ* conservation of *Cymbidium finlaysonianum* by seed storage. *Biodiversitas*, 21(8), 3519–3524. <https://doi.org/10.13057/biodiv/d210813>
- Ramírez-Mosqueda, M. A., Cruz-Cruz, C. A., Atlahua-Temoxtle, J., & Bello-Bello, J. J. (2019). *In vitro* conservation and regeneration of *Laelia anceps* Lindl. *South African Journal of Botany*, 121, 219–223. <https://doi.org/10.1016/j.sajb.2018.11.010>
- Ruta, C., Lombardi, M., & Ozudogru, E. A. (2020). Biobanking of vegetable genetic resources by *in vitro* conservation and cryopreservation. In *Biodiversity and Conservation* 29(13), 3495–3532. Springer Science and Business Media B.V. <https://doi.org/10.1007/s10531-020-02051-0>
- Salazar-Mercado, S. A., & Botello-Delgado, E. A. (2020). Effect of the medium composition on the asymbiotic germination and *in vitro* development of the *Laeliocattleya* hybrid. *South African Journal of Botany*, 135, 80–86. <https://doi.org/10.1016/j.sajb.2020.08.011>
- Sanghamitra, P., Samantaray, S., Bagchi, T. B., & Mandal, B. B. (2019). Conservation of medicinal yam *in vitro*: Effect of ionic strength, sucrose, mannitol, ABA and low temperature. *Indian Journal of Horticulture*, 76(4), 701–706. <https://doi.org/10.5958/0974-0112.2019.00110.5>
- Santos Lima, A. P., Oliveira Bastos, F. de J., Lima Brito, A., Borges Fernandes Gilenio, & Ferreira de Santana, J. de R. (2021). Modulation of culture medium on the *ex situ* conservation of *Neoregelia mucugensis* Leme (Bromeliaceae). *Revista Caatinga*, 34(4), 763–771.
- Seaton, P., Kendon, J., Pritchard, H. W., Puspitaningtyas, M., & Marks, T. R. (2013). Orchid conservation: the next ten years. *Lankesteriana International Journal on Orchidology*, 13(2).
- Sharma, J., Koul, A., Sharma, S., Shankarayyan, R., & Mallubhotla, S. (2021). *In vitro* propagation of *Nanorrhynchus ramosissimum* (Wall.) Betsche: A traditionally important medicinal plant.

Recibido:
3/agosto/2021

Aceptado:
12/enero/2022

- Sianipar, N. F., & Mariska, I. (2020). Micropropagation of *Orthosiphon aristatus* through indirect and direct organogenesis. *Jurnal Teknologi*, 82(3), 45–51. <https://doi.org/10.11113/jt.v82.13887>
- Singh, A., Shukla, P. K., Sengar, R. S., & Mishra, P. (2021). *In vitro* effect of polyethylene glycol and sorbitol on two banana varieties viz. *Grand naine* and *Nalla bontha* to study drought stress. *Journal of Applied and Natural Science*, 13(2), 482–490. <https://doi.org/10.31018/jans.v13i2.2579>
- Sorgato, J. C., Soares, J. S., Damiani, C. R., & Ribeiro, L. M. (2020). Effects of light, agar, activated charcoal, and culture medium on the germination and early development of *dendrobium* seedlings. *Australian Journal of Crop Science*, 14(4), 557–564. <https://doi.org/10.21475/ajcs.20.14.04.p1528>
- Tavazza, R., Rey, N. A., & Pagnotta M.A. (2015). Globe artichoke *in vitro* conservation protocol to meet germplasm preservation and production management. *IX International Symposium on Artichoke, Cardoon and Their Wild Relatives*, 421–428.
- Than, M. M. M. (2013). *In vitro* conservation of endangered orchid *Bulbophyllum auricomum* lindl., the royal orchid of Myanmar. *Propagation of Ornamental Plants*, 13(4), 154–159.
- Tokuhara, K., Mii, M., & Phillips, G. C. (1993). Micropropagation of *Phalaenopsis* and *Doritaenopsis* by culturing shoot tips of flower stalk buds. In *Plant Cell Reports* (Vol. 13).
- Utami, E. S. W., & Hariyanto, S. (2019). *In vitro* seed germination and seedling development of a rare indonesian native orchid *Phalaenopsis amboinensis* J.J.Sm. *Scientifica*, 2019. <https://doi.org/10.1155/2019/8105138>
- Velázquez Kú, V. N., Quijano-Ávila, J. del C., & Rodríguez-Ávila, N.L. (2016). Análisis de diferentes sustratos en la germinación y multiplicación *in vitro* de orquídeas silvestres del estado de Campeche. *Revista Del Centro de Graduados e Investigación, Instituto Tecnológico de Mérida*, 27–31.
- Vudala, S. M., & Ribas, L. L. F. (2017). Seed storage and asymbiotic germination of *Hadrolaelia grandis* (Orchidaceae). *South African Journal of Botany*, 108, 1–7. <https://doi.org/10.1016/j.sajb.2016.09.008>
- Yam, T. W., & Arditti, J. (2009). History of orchid propagation: A mirror of the history of biotechnology. In *Plant Biotechnology Reports* 3(1), 1–56. <https://doi.org/10.1007/s11816-008-0066-3>
- Yamazaki, J., & Miyoshi, K. (2006). *In vitro* asymbiotic germination of immature seed and formation of protocorm by *Cephalanthera falcata* (Orchidaceae). *Annals of Botany*, 98(6), 1197–1206. <https://doi.org/10.1093/aob/mcl223>
- Yeh, C. H., Chen, K. Y., & Lee, Y. I. (2021). Asymbiotic germination of *Vanilla planifolia* in relation to the timing of seed collection and seed pretreatments. *Botanical Studies*, 62(1). <https://doi.org/10.1186/s40529-021-00311-y>